

4-28-00

A

UTILITY PATENT APPLICATION TRANSMITTAL <i>(Only for new nonprovisional applications under 37 CFR 1.53(b))</i>	Attorney Docket No.	L0461/7086 (JRV)
	First Named Inventor or Application Identifier	
	Ono et al.	
	Express Mail Label No.	EL018097968US
	Date of Deposit	April 26, 2000

09/559013
 04/27/00

APPLICATION ELEMENTS <i>See MPEP chapter 600 concerning utility patent application contents</i>	ADDRESS TO: Box Patent Application Assistant Commissioner for Patents Washington, DC 20231
1. <input type="checkbox"/> Fee Transmittal Form <i>(Submit an original, and a duplicate for fee processing)</i> 2. <input checked="" type="checkbox"/> Application [Total pages 80] 64 - pages specification 1 - pages abstract 15 - pages claims 121 - Total claims 3. <input type="checkbox"/> Drawing(s) (35 USC 113) [Total sheets] <input type="checkbox"/> Informal <input type="checkbox"/> Formal [Total drawings] 4. <input type="checkbox"/> Oath or Declaration [Total pages] a. <input type="checkbox"/> Newly executed (original or copy) b. <input type="checkbox"/> Unsigned c. <input type="checkbox"/> Copy from a prior application (37 CFR 1.63(d)) <i>(for continuation/divisional with Box 17 completed)</i> [Note Box 5 below] i. <input type="checkbox"/> DELETION OF INVENTOR(S) Signed statement attached deleting inventor(s) named in the prior application, see 37 CFR 1.63(d)(2) and 1.33(b). 5. <input type="checkbox"/> Incorporation by Reference <i>(usable if Box 4c is checked)</i> The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied under Box 4c, is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein.	6. <input type="checkbox"/> Microfiche Computer Program (Appendix) 7. <input checked="" type="checkbox"/> Nucleotide and/or Amino Acid Sequence Submission (if applicable, all necessary) a. <input checked="" type="checkbox"/> Computer Readable Copy b. <input checked="" type="checkbox"/> Paper Copy (identical to computer copy) c. <input checked="" type="checkbox"/> Statement verifying identity of above copies ACCOMPANYING APPLICATION PARTS 8. <input type="checkbox"/> Assignment Papers/cover sheet & documents(s) 9. <input type="checkbox"/> 37 CFR 3.73(b) Statement <i>(when there is an assignee)</i> <input type="checkbox"/> Power of Attorney 10. <input type="checkbox"/> English Translation of Document <i>(if applicable)</i> 11. <input type="checkbox"/> Information Disclosure Statement PTO-1449 <input type="checkbox"/> Copies of IDS Citations 12. <input type="checkbox"/> Preliminary Amendment 13. <input checked="" type="checkbox"/> Return Receipt Postcard (MPEP 503) <i>(Should be specifically itemized)</i> 14. <input type="checkbox"/> Small Entity Statement(s) <input type="checkbox"/> Statement filed in prior application, Status still proper and desired 15. <input type="checkbox"/> Certified Copy of Priority Document(s) <i>(if foreign priority is claimed)</i>
16. Other:	
17. If a CONTINUING APPLICATION , check appropriate box and supply the requisite information: <input type="checkbox"/> Continuation <input type="checkbox"/> Divisional <input type="checkbox"/> Continuation-in-part (CIP) of prior application No.: <input type="checkbox"/> Cancel in this application original claims of the prior application before calculating the filing fee. <input type="checkbox"/> Amend the specification by inserting before the first line the sentence: This application is a <input type="checkbox"/> continuation <input type="checkbox"/> divisional of application serial no. , filed , entitled , and now .	

18. CORRESPONDENCE ADDRESS					
Correspondence address below					
ATTORNEY'S NAME	John R. Van Amsterdam, Reg. No. 40,212				
NAME	Wolf, Greenfield & Sacks, P.C.				
ADDRESS	600 Atlantic Avenue				
CITY	Boston	STATE	MA	ZIP	02210
COUNTRY	USA	TELEPHONE	(617) 720-3500	FAX	(617) 720-2441

19. SIGNATURE OF APPLICANT, ATTORNEY, OR AGENT REQUIRED	
NAME	John R. Van Amsterdam, Reg. No. 40,212
SIGNATURE	<i>John R. Van Amsterdam</i>
DATE	APRIL 26, 2000

CANCER ASSOCIATED ANTIGENS AND USES THEREFOR

Related Applications

This application claims priority under 35 U.S.C. §119 from U.S. provisional
5 application serial number 60/168,353, filed December 1, 1999.

Field of the Invention

The invention relates to nucleic acids and encoded polypeptides which are cancer
associated antigens expressed in methylcholanthrene-induced fibrosarcomas. The invention
10 also relates to agents which bind the nucleic acids or polypeptides. The nucleic acid
molecules, polypeptides coded for by such molecules and peptides derived therefrom, as well
as related antibodies and cytolytic T lymphocytes, are useful, *inter alia*, in diagnostic and
therapeutic contexts.

Background of the Invention

The mechanism by which T cells recognize foreign materials has been implicated in
cancer. A number of cytolytic T lymphocyte (CTL) clones directed against autologous
melanoma antigens, testicular antigens, and melanocyte differentiation antigens have been
described. In many instances, the antigens recognized by these clones have been
20 characterized.

The use of autologous CTLs for identifying tumor antigens requires that the target
cells which express the antigens can be cultured *in vitro* and that stable lines of autologous
CTL clones which recognize the antigen-expressing cells can be isolated and propagated.
While this approach has worked well for melanoma antigens, other tumor types, such as
25 epithelial cancers including breast and colon cancer, have proved refractory to the approach.

More recently another approach to the problem has been described by Sahin et al.
(*Proc. Natl. Acad. Sci. USA* 92:11810-11813, 1995). According to this approach, autologous
antisera are used to identify immunogenic protein antigens expressed in cancer cells by
screening expression libraries constructed from tumor cell cDNA. Antigen-encoding clones
30 so identified have been found to have elicited an high-titer humoral immune response in the
patients from which the antisera were obtained. Such a high-titer IgG response implies helper
T cell recognition of the detected antigen. These tumor antigens can then be screened for the
presence of MHC/HLA class I and class II motifs and reactivity with CTLs.

Presently there is a need for additional cancer antigens for development of therapeutics and diagnosis applicable to a greater number of cancer patients having various cancers.

Summary of the Invention

5 Autologous antibody screening has now been applied to methylcholanthrene-induced fibrosarcomas using antisera from mice bearing such tumors. Numerous cancer associated antigens have been identified, including human nucleic acid homologs of mouse clones so identified. The invention provides, *inter alia*, isolated nucleic acid molecules, expression vectors containing those molecules and host cells transfected with those molecules. The
10 invention also provides isolated proteins and peptides, antibodies to those proteins and peptides and CTLs which recognize the proteins and peptides. Fragments including functional fragments and variants of the foregoing also are provided. Kits containing the foregoing molecules additionally are provided. The foregoing can be used in the diagnosis, monitoring, research, or treatment of conditions characterized by the expression of one or
15 more cancer associated antigens.

The invention involves the surprising discovery of several genes, some previously known and some previously unknown, which are expressed in mice bearing methylcholanthrene-induced fibrosarcomas. These individuals all have serum antibodies against the proteins (or fragments thereof) encoded by these genes. Thus, abnormally
20 expressed genes are recognized by the host's immune system and therefore can form a basis for diagnosis, monitoring and therapy. The human equivalents (i.e., homologs) of the nucleic acids have in some cases also been identified.

The invention involves the use of a single material, a plurality of different materials and even large panels and combinations of materials. For example, a single gene, a single
25 protein encoded by a gene, a single functional fragment thereof, a single antibody thereto, etc. can be used in methods and products of the invention. Likewise, pairs, groups and even panels of these materials and optionally other cancer associated antigen genes and/or gene products can be used for diagnosis, monitoring and therapy. The pairs, groups or panels can involve 2, 3, 4, 5 or more genes, gene products, fragments thereof or agents that recognize
30 such materials. A plurality of such materials are not only useful in monitoring, typing, characterizing and diagnosing cells abnormally expressing such genes, but a plurality of such materials can be used therapeutically. An example of the use of a plurality of such materials for the prevention, delay of onset, amelioration, etc. of cancer cells, which express or will

express such genes prophylactically or acutely. Any and all combinations of the genes, gene products, and materials which recognize the genes and gene products can be tested and identified for use according to the invention. It would be far too lengthy to recite all such combinations; those skilled in the art, particularly in view of the teaching contained herein, will readily be able to determine which combinations are most appropriate for which circumstances.

As will be clear from the following discussion, the invention has *in vivo* and *in vitro* uses, including for therapeutic, diagnostic, monitoring and research purposes. One aspect of the invention is the ability to fingerprint a cell expressing a number of the genes identified according to the invention by, for example, quantifying the expression of such gene products. Such fingerprints will be characteristic, for example, of the stage of the cancer, the type of the cancer, or even the effect in animal models of a therapy on a cancer. Cells also can be screened to determine whether such cells abnormally express the genes identified according to the invention.

The invention, in one aspect, is a method of diagnosing a disorder characterized by expression of a cancer associated antigen precursor coded for by a nucleic acid molecule. The method involves the steps of contacting a biological sample isolated from a subject with an agent that specifically binds to the nucleic acid molecule, an expression product thereof, or a fragment of an expression product thereof complexed with a MHC, preferably an HLA, molecule, wherein the nucleic acid molecule is a NA Group 1 nucleic acid molecule, and determining the interaction between the agent and the nucleic acid molecule, the expression product or fragment of the expression product as a determination of the disorder.

In one embodiment the agent is selected from the group consisting of (a) a nucleic acid molecule comprising NA Group 1 nucleic acid molecules or a fragment thereof, (b) a nucleic acid molecule comprising NA Group 3 nucleic acid molecules or a fragment thereof, (c) a nucleic acid molecule comprising NA Group 5 nucleic acid molecules or a fragment thereof, (d) an antibody that binds to an expression product, or a fragment thereof, of NA group 1 nucleic acids, (e) an antibody that binds to an expression product, or a fragment thereof, of NA group 3 nucleic acids, (f) an antibody that binds to an expression product, or a fragment thereof, of NA group 5 nucleic acids, (g) and agent that binds to a complex of an MHC, preferably HLA, molecule and a fragment of an expression product of a NA Group 1 nucleic acid, (h) an agent that binds to a complex of an MHC, preferably HLA, molecule and a fragment of an expression product of a NA group 3 nucleic acid, and (i) an agent that binds

to a complex of an MHC, preferably HLA, molecule and a fragment of an expression product of a NA Group 5 nucleic acid.

The disorder may be characterized by expression of a plurality of cancer associated antigen precursors. Thus the methods of diagnosis may include use of a plurality of agents, each of which is specific for a different cancer associated antigen precursor (including at least one of the cancer associated antigen precursors disclosed herein), and wherein said plurality of agents is at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9 or at least 10 such agents.

In each of the above embodiments the agent may be specific for a cancer associated antigen precursor, including the cancer associated antigen precursors disclosed herein, and preferably human homologs thereof.

In another aspect the invention is a method for determining regression, progression or onset of a condition characterized by expression of abnormal levels of a protein encoded by a nucleic acid molecule that is a NA Group 1 molecule. The method involves the steps of monitoring a sample, from a subject who has or is suspected of having the condition, for a parameter selected from the group consisting of (i) the protein, (ii) a peptide derived from the protein, (iii) an antibody which selectively binds the protein or peptide, and (iv) cytolytic T cells specific for a complex of the peptide derived from the protein and a MHC molecule, as a determination of regression, progression or onset of said condition. In one embodiment the sample is a body fluid, a body effusion or a tissue.

In another embodiment the step of monitoring comprises contacting the sample with a detectable agent selected from the group consisting of (a) an antibody which selectively binds the protein of (i), or the peptide of (ii), (b) a protein or peptide which binds the antibody of (iii), and (c) a cell which presents the complex of the peptide and MHC molecule of (iv). In a preferred embodiment the antibody, the protein, the peptide or the cell is labeled with a radioactive label or an enzyme. The sample in a preferred embodiment is assayed for the peptide.

According to another embodiment the nucleic acid molecule is one of the following: a NA Group 3 molecule or a NA Group 5 molecule. In yet another embodiment the protein is a plurality of proteins, the parameter is a plurality of parameters, each of the plurality of parameters being specific for a different of the plurality of proteins.

The invention in another aspect is a pharmaceutical preparation, preferably for a human subject. The pharmaceutical preparation includes an agent which when administered

to the subject enriches selectively the presence of complexes of a MHC molecule, preferably a HLA molecule, and a cancer associated antigen, and a pharmaceutically acceptable carrier, wherein the cancer associated antigen is a fragment of a cancer associated antigen precursor encoded by a nucleic acid molecule which comprises a NA Group 1 molecule, or a human homolog thereof. In one embodiment the nucleic acid molecule is a NA Group 3 nucleic acid molecule.

The agent in one embodiment comprises a plurality of agents, each of which enriches selectively in the subject complexes of a MHC molecule and a different cancer associated antigen. Preferably the plurality is at least two, at least three, at least four or at least 5 different such agents.

In another embodiment the agent is selected from the group consisting of (1) an isolated polypeptide comprising the cancer associated antigen, a human homolog thereof, or a functional variant thereof, (2) an isolated nucleic acid operably linked to a promoter for expressing the isolated polypeptide, homolog or functional variant thereof, (3) a host cell expressing the isolated polypeptide, homolog or functional variant thereof, and (4) isolated complexes of the polypeptide, homolog or functional variants thereof, and an HLA molecule.

The agent may be a cell expressing an isolated polypeptide. In one embodiment the agent is a cell expressing an isolated polypeptide comprising the cancer associated antigen, a human homolog thereof or a functional variant thereof. In another embodiment the agent is a cell expressing an isolated polypeptide comprising the cancer associated antigen, a homolog or a functional variant thereof, and wherein the cell expresses an HLA molecule that binds the polypeptide. The cell can express one or both of the polypeptide and HLA molecule recombinantly. In preferred embodiments the cell is nonproliferative. In yet another embodiment the agent is at least two, at least three, at least four or at least five different polypeptides, each representing a different cancer associated antigen, homolog or functional variant thereof.

The agent in one embodiment is a PP Group 2 polypeptide. In other embodiments the agent is a PP Group 3 polypeptide or a PP Group 4 polypeptide.

In an embodiment each of the pharmaceutical preparations described herein also includes an adjuvant.

According to another aspect the invention, a composition is provided which includes an isolated agent that binds selectively a PP Group 1 polypeptide. In separate embodiments the agent binds selectively to a polypeptide selected from the following: a PP Group 2

polypeptide, a PP Group 3 polypeptide, a PP Group 4 polypeptide, and a PP Group 5 polypeptide. In other embodiments, the agent is a plurality of different agents that bind selectively at least two, at least three, at least four, or at least five different such polypeptides. In each of the above described embodiments the agent may be an antibody.

5 In another aspect the invention is a composition of matter composed of a conjugate of the agent of the above-described compositions of the invention and a therapeutic or diagnostic agent. Preferably the conjugate is of the agent and a therapeutic or diagnostic that is an antineoplastic.

10 The invention in another aspect is a pharmaceutical composition which includes an isolated nucleic acid molecule selected from the group consisting of: (1) NA Group 1 molecules, and (2) NA Group 2 molecules, and a pharmaceutically acceptable carrier. In one embodiment the isolated nucleic acid molecule comprises a NA Group 3 or NA Group 4 molecule. In another embodiment the isolated nucleic acid molecule comprises at least two isolated nucleic acid molecules coding for two different polypeptides, each polypeptide
15 comprising a different cancer associated antigen.

Preferably the pharmaceutical composition also includes an expression vector with a promoter operably linked to the isolated nucleic acid molecule. In another embodiment the pharmaceutical composition also includes a host cell recombinantly expressing the isolated nucleic acid molecule.

20 According to another aspect of the invention a pharmaceutical composition is provided. The pharmaceutical composition includes an isolated polypeptide comprising a PP Group 1 or a PP Group 2 polypeptide, and a pharmaceutically acceptable carrier. In one embodiment the isolated polypeptide comprises a PP Group 3 or a PP Group 4 polypeptide.

In another embodiment the isolated polypeptide comprises at least two different
25 polypeptides, each comprising a different cancer associated antigen at least one of which is encoded by a NA group 1 molecule as disclosed herein. In separate embodiments the isolated polypeptides are selected from the following: PP Group 3 polypeptides or HLA binding fragments thereof and PP Group 5 polypeptides or HLA binding fragments thereof.

In an embodiment each of the pharmaceutical compositions described herein also
30 includes an adjuvant.

Another aspect the invention is an isolated nucleic acid molecule comprising a NA Group 3 molecule. Another aspect the invention is an isolated nucleic acid molecule comprising a NA Group 4 molecule.

The invention in another aspect is an isolated nucleic acid molecule selected from the group consisting of (a) a fragment of a nucleic acid selected from the group of nucleic acid molecules consisting of SEQ ID NOs numbered below and comprising all nucleic acid sequences among SEQ ID Nos: 9, 13, 15, 17, 19 and 23, of sufficient length to represent a sequence unique within the mouse or human genomes, and identifying a nucleic acid encoding a cancer associated antigen precursor, (b) complements of (a), provided that the fragment includes a sequence of contiguous nucleotides which is not identical to any sequence selected from the sequence group consisting of (1) sequences having the GenBank accession numbers of Table 8, (2) complements of (1), and (3) fragments of (1) and (2).

In one embodiment the sequence of contiguous nucleotides is selected from the group consisting of: (1) at least two contiguous nucleotides nonidentical to the sequences in Table 8, (2) at least three contiguous nucleotides nonidentical to the sequences in Table 8, (3) at least four contiguous nucleotides nonidentical to the sequences in Table 8, (4) at least five contiguous nucleotides nonidentical to the sequences in Table 8, (5) at least six contiguous nucleotides nonidentical to the sequences in Table 8, or (6) at least seven contiguous nucleotides nonidentical to the sequences in Table 8.

In another embodiment the fragment has a size selected from the group consisting of at least: 8 nucleotides, 10 nucleotides, 12 nucleotides, 14 nucleotides, 16 nucleotides, 18 nucleotides, 20, nucleotides, 22 nucleotides, 24 nucleotides, 26 nucleotides, 28 nucleotides, 30 nucleotides, 50 nucleotides, 75 nucleotides, 100 nucleotides, 200 nucleotides, 1000 nucleotides and every integer length therebetween.

In yet another embodiment the molecule encodes a polypeptide which, or a fragment of which, binds a MHC molecule, preferably a human HLA receptor, or an antibody, preferably one having human amino acid sequences.

Another aspect of the invention is an expression vector comprising an isolated nucleic acid molecule of the invention described above operably linked to a promoter.

According to one aspect the invention is an expression vector comprising a nucleic acid operably linked to a promoter, wherein the nucleic acid is a NA Group 1 or Group 2 molecule. In another aspect the invention is an expression vector comprising a NA Group 1 or Group 2 molecule and a nucleic acid encoding an MHC, preferably HLA, molecule.

In yet another aspect the invention is a host cell transformed or transfected with an expression vector of the invention described above.

In another aspect the invention is a host cell transformed or transfected with an

expression vector comprising an isolated nucleic acid molecule of the invention described above operably linked to a promoter, or an expression vector comprising a nucleic acid operably linked to a promoter, wherein the nucleic acid is a NA Group 1 or 2 molecule and further comprising a nucleic acid encoding a MHC molecule.

5 According to another aspect of the invention an isolated polypeptide encoded by the isolated nucleic acid molecules the invention, described above, is provided. These include PP Group 1-5 polypeptides, including SEQ ID NOS:2, 4, 6 and 8. The invention also includes a fragment of the polypeptide which is immunogenic. In one embodiment the fragment, or a portion of the fragment, binds a MHC molecule, preferably HLA ,or an antibody, preferably
10 one having human amino acid sequences.

 The invention includes in another aspect an isolated fragment of a cancer associated antigen precursor which, or portion of which, binds a MHC molecule, preferably HLA, or a human antibody, or an antibody, preferably one having human amino acid sequences, wherein the precursor is encoded by a nucleic acid molecule that is a NA Group 1 molecule. In one
15 embodiment the fragment is part of a complex with MHC/HLA. In another embodiment the fragment is between 8 and 12 amino acids in length. In another embodiment the invention includes an isolated polypeptide comprising a fragment of the polypeptide of sufficient length to represent a sequence unique within the mouse or human genomes and identifying a polypeptide that is a cancer associated antigen precursor.

20 According to another aspect of the invention a kit for detecting the presence of the expression of a cancer associated antigen precursor is provided. The kit includes a pair of isolated nucleic acid molecules each of which consists essentially of a molecule selected from the group consisting of (a) a 12-32 nucleotide contiguous segment of the nucleotide sequence of any of the NA Group 1 molecules and (b) complements of (“a”), wherein the contiguous
25 segments are nonoverlapping. In one embodiment the pair of isolated nucleic acid molecules is constructed and arranged to selectively amplify an isolated nucleic acid molecule that is a NA Group 3 molecule. Preferably, the pair amplifies a human NA Group 3 molecule.

 According to another aspect of the invention a method for treating a subject with a disorder characterized by expression of a cancer associated antigen precursor is provided. The
30 method includes the step of administering to the subject an amount of an agent, which enriches selectively in the subject the presence of complexes of a MHC/HLA molecule and a cancer associated antigen, effective to ameliorate the disorder, wherein the cancer associated antigen is a fragment of a cancer associated antigen precursor encoded by a nucleic acid

molecule selected from the group consisting of (a) a nucleic acid molecule comprising NA group 1 nucleic acid molecules, (b) a nucleic acid molecule comprising NA group 3 nucleic acid molecules, (c) a nucleic acid molecule comprising NA group 5 nucleic acid molecules.

5 In one embodiment the disorder is characterized by expression of a plurality of cancer associated antigen precursors and wherein the agent is a plurality of agents, each of which enriches selectively in the subject the presence of complexes of a MHC/HLA molecule and a different cancer associated antigen. Preferably the plurality is at least 2, at least 3, at least 4, or at least 5 such agents.

10 In another embodiment the agent is an isolated polypeptide selected from the group consisting of PP Group 1, PP Group 2, PP Group 3, PP Group 4, and PP group 5 polypeptides.

In yet another embodiment the disorder is cancer.

15 According to another aspect the invention is a method for treating a subject having a condition characterized by expression of a cancer associated antigen precursor in cells of the subject. The method includes the steps of (i) removing an immunoreactive cell containing sample from the subject, (ii) contacting the immunoreactive cell containing sample to the host cell under conditions favoring production of cytolytic T cells against a cancer associated antigen which is a fragment of the precursor, (iii) introducing the cytolytic T cells to the subject in an amount effective to lyse cells which express the cancer associated antigen,
20 wherein the host cell is transformed or transfected with an expression vector comprising an isolated nucleic acid molecule operably linked to a promoter, the isolated nucleic acid molecule being selected from the group of nucleic acid molecules consisting of NA Group 1, NA Group 2, NA Group 3, NA Group 4, NA Group 5.

25 In one embodiment the host cell recombinantly expresses an HLA molecule which binds the cancer associated antigen. In another embodiment the host cell endogenously expresses a MHC/HLA molecule which binds the cancer associated antigen.

The invention includes in another aspect a method for treating a subject having a condition characterized by expression of a cancer associated antigen precursor in cells of the subject. The method includes the steps of (i) identifying a nucleic acid molecule expressed by
30 the cells associated with said condition, wherein said nucleic acid molecule is a NA Group 1 molecule (ii) transfecting a host cell with a nucleic acid selected from the group consisting of (a) the nucleic acid molecule identified, (b) a fragment of the nucleic acid identified which includes a segment coding for a cancer associated antigen, (c) deletions, substitutions or

additions to (a) or (b), and (d) degenerates of (a), (b), or (c); (iii) culturing said transfected host cells to express the transfected nucleic acid molecule, and; (iv) introducing an amount of said host cells or an extract thereof to the subject effective to increase an immune response against the cells of the subject associated with the condition. Preferably, the antigen is a human antigen and the subject is a human.

In one embodiment the method also includes the step of (a) identifying a MHC molecule which presents a portion of an expression product of the nucleic acid molecule, wherein the host cell expresses the same MHC molecule as identified in (a) and wherein the host cell presents a MHC binding portion of the expression product of the nucleic acid molecule.

In another embodiment the method also includes the step of treating the host cells to render them non-proliferative.

In yet another embodiment the immune response comprises a B-cell response or a T cell response. Preferably the response is a T-cell response which comprises generation of cytolytic T-cells specific for the host cells presenting the portion of the expression product of the nucleic acid molecule or cells of the subject expressing the cancer associated antigen.

In another embodiment the nucleic acid molecule is a NA Group 3 molecule.

Another aspect of the invention is a method for treating or diagnosing or monitoring a subject having a condition characterized by expression of an abnormal amount of a protein encoded by a nucleic acid molecule that is a NA Group 1 molecule. The method includes the step of administering to the subject an antibody which specifically binds to the protein or a peptide derived therefrom, the antibody being coupled to a therapeutically useful agent, in an amount effective to treat the condition.

In one embodiment the antibody is a monoclonal antibody. Preferably the monoclonal antibody is a chimeric antibody or a humanized antibody.

In another aspect the invention is a method for treating a condition characterized by expression in a subject of abnormal amounts of a protein encoded by a nucleic acid molecule that is a NA Group 1 nucleic acid molecule. The method involves the step of administering to a subject at least one of the pharmaceutical compositions of the invention described above in an amount effective to prevent, delay the onset of, or inhibit the condition in the subject. In one embodiment the condition is cancer. In another embodiment the method includes the step of first identifying that the subject expresses in a tissue abnormal amounts of the protein.

The invention in another aspect is a method for treating a subject having a condition

characterized by expression of abnormal amounts of a protein encoded by a nucleic acid molecule that is a NA Group 1 nucleic acid molecule. The method includes the steps of (i) identifying cells from the subject which express abnormal amounts of the protein; (ii) isolating a sample of the cells; (iii) cultivating the cells, and (iv) introducing the cells to the
5 subject in an amount effective to provoke an immune response against the cells.

In one embodiment the method includes the step of rendering the cells non-proliferative, prior to introducing them to the subject.

In another aspect the invention is a method for treating a pathological cell condition characterized by abnormal expression of a protein encoded by a nucleic acid molecule that is a
10 NA Group 1 nucleic acid molecule. The method includes the step of administering to a subject in need thereof an effective amount of an agent which inhibits the expression or activity of the protein.

In one embodiment the agent is an inhibiting antibody which selectively binds to the protein and wherein the antibody is a monoclonal antibody, a chimeric antibody, a humanized
15 antibody or a fragment thereof. In another embodiment the agent is an antisense nucleic acid molecule which selectively binds to the nucleic acid molecule which encodes the protein. In yet another important embodiment the nucleic acid molecule is a NA Group 3 nucleic acid molecule.

The invention includes in another aspect a composition of matter useful in stimulating
20 an immune response to a plurality of proteins encoded by nucleic acid molecules that are NA Group 1 molecules. The composition is a plurality of peptides derived from the amino acid sequences of the proteins, wherein the peptides bind to one or more MHC molecules presented on the surface of the cells which express an abnormal amount of the protein.

In one embodiment at least a portion of the plurality of peptides bind to MHC
25 molecules and elicit a cytolytic response thereto. In another embodiment the composition of matter includes an adjuvant. In another embodiment the adjuvant is a saponin, GM-CSF, or an interleukin. In still another embodiment, the compositions also includes at least one peptide useful in stimulating an immune response to at least one protein which is not encoded by nucleic acid molecules that are NA Group 1 molecules, wherein the at least one peptide
30 binds to one or more MHC molecules.

According to another aspect the invention is an isolated antibody which selectively binds to a complex of: (i) a peptide derived from a protein encoded by a nucleic acid molecule that is a NA Group 1 molecule and (ii) and an MHC molecule to which binds the peptide to

form the complex, wherein the isolated antibody does not bind to (i) or (ii) alone.

In one embodiment the antibody is a monoclonal antibody, a chimeric antibody, a humanized antibody or a fragment thereof.

The invention also involves the use of the genes, gene products, fragments thereof, agents which bind thereto, and so on in the preparation of medicaments. A particular medicament is for treating cancer, preferably bladder cancer, colon cancer, lung cancer, breast cancer or hepatoma.

In each of the foregoing embodiments, a preferred nucleic acid molecule include the nucleotide sequence of SEQ ID NO:23 or fragments thereof, and polypeptides comprise SEQ ID NO:24 or fragments thereof, or are encoded by SEQ ID NO:23 or fragments thereof.

These and other aspects of the invention will be described in further detail in connection with the detailed description of the invention.

Detailed Description of the Invention

In the above summary and in the ensuing description, lists of sequences are provided. The lists are meant to embrace each single sequence separately, two or more sequences together where they form a part of the same gene, any combination of two or more sequences which relate to different genes, including and up to the total number on the list, as if each and every combination were separately and specifically enumerated. Likewise, when mentioning fragment size, it is intended that a range embrace the smallest fragment mentioned to the full-length of the sequence (less one nucleotide or amino acid so that it is a fragment), each and every fragment length intended as if specifically enumerated. Thus, if a fragment could be between 10 and 15 in length, it is explicitly meant to mean 10, 11, 12, 13, 14, or 15 in length.

The summary and the claims mention antigen precursors and antigens. As used in the summary and in the claims, a precursor is substantially the full-length protein encoded by the coding region of the isolated DNA and the antigen is a peptide which complexes with MHC, preferably HLA, and which participates in the immune response as part of that complex. Such antigens are typically 9 amino acids long, although this may vary slightly.

As used herein, a subject is a human, non-human primate, cow, horse, pig, sheep, goat, dog, cat or rodent. In all embodiments human cancer associated antigens, which have substantial nucleotide and/or amino acid sequence identity to the presently identified cancer associated antigens, and human subjects are preferred.

The present invention in one aspect involves the cloning of cDNAs encoding cancer

associated antigen precursors using autologous antisera of mice having methylcholanthrene-induced fibrosarcomas. The sequences of the clones representing genes identified according to the methods described herein are presented in the attached Sequence Listing. Of the foregoing, it can be seen that some of the clones are considered completely novel as no
5 nucleotide or amino acid homologies to coding regions were found in the databases searched. Other clones are novel but have some homology to sequences deposited in databases (mainly EST sequences). Nevertheless, the entire gene sequence was not previously known. In some cases no function was suspected and in other cases, even if a function was suspected, it was not known that the gene was associated with cancer. In all cases, it was not known or
10 suspected that the gene encoded a cancer antigen which reacted with antibody from autologous sera. Analysis of the clone sequences by comparison to nucleic acid and protein databases determined that still other of the clones surprisingly are closely related to other previously-cloned genes. The sequences of these related genes is also presented in the Sequence Listing. The nature of the foregoing genes as encoding antigens recognized by the
15 immune systems of cancer patients is, of course, unexpected.

The invention thus involves in one aspect cancer associated antigen polypeptides, genes encoding those polypeptides, functional modifications and variants of the foregoing, useful fragments of the foregoing, as well as diagnostics and therapeutics relating thereto.

Homologs and alleles of the cancer associated antigen nucleic acids of the invention
20 can be identified by conventional techniques. Thus, an aspect of the invention is those nucleic acid sequences which code for cancer associated antigen precursors. The following chart is provided to identify the various groups of sequences discussed in the claims and in the summary:

25 Nucleic Acid Sequences

NA Group 1. (a) nucleic acid molecules which hybridize under stringent conditions to a molecule consisting of a nucleic acid sequence selected from the group consisting of nucleic acid sequences among SEQ ID NOs: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19 and 23 and which code for
30 a cancer associated antigen precursor,

(b) deletions, additions and substitutions which code for a respective cancer associated antigen precursor,

(c) nucleic acid molecules that differ from the nucleic acid molecules of (a) or

(b) in codon sequence due to the degeneracy of the genetic code, and

(d) complements of (a), (b) or (c).

NA Group 2. Fragments of NA Group 1, which codes for a polypeptide which, or a portion
5 of which, binds an MHC molecule to form a complex recognized by an autologous antibody
or lymphocyte.

NA Group 3. The subset of NA Group 1 where the nucleotide sequence is selected from the
group consisting of:

10 (a) previously unknown nucleic acids coding for a cancer associated antigen
precursor,

(b) deletions, additions and substitutions which code for a respective cancer
associated antigen precursor,

(c) nucleic acid molecules that differ from the nucleic acid molecules of (a) or

15 (b) in codon sequence due to the degeneracy of the genetic code, and

(d) complements of (a), (b) or (c).

NA Group 4. Fragments of NA Group 3, which code for a polypeptide which, or a portion of
20 which, binds to an MHC molecule to form a complex recognized by an autologous antibody
or lymphocyte.

NA Group 5. A subset of NA Group 1, comprising cancer associated antigens that react with
allogeneic cancer antisera.

25 Polypeptide Sequences

PP Group 1. Polypeptides encoded by NA Group 1.

PP Group 2. Polypeptides encoded by NA Group 2

PP Group 3. Polypeptides encoded by NA Group 3.

PP Group 4. Polypeptides encoded by NA Group 4.

30 PP Group 5. Polypeptides encoded by NA Group 5.

Identification of human homologs of the cancer associated antigens will be familiar to
those of skill in the art. In particular, the methods described in Example 8 were effective in

identifying the human homolog of OY-MC-4, OY-TES-1, as well as establishing it as a cancer-testis antigen useful in therapeutic and diagnostic applications for treatment and diagnosis of cancer. In general, nucleic acid hybridization is a suitable method for identification of homologous sequences of another species (e.g., human) which correspond to a known sequence (e.g., mouse cancer associated sequences presented herein). Standard nucleic acid hybridization procedures can be used to identify related nucleic acid sequences of selected precent identity. For example, one can construct a library of cDNAs reverse transcribed from the mRNA of a selected tissue (e.g., testis) and use the cancer associated antigen nucleic acids identified herein to screen the library for related nucleotide sequences. The screening can be performed at various stringencies to identify those sequences which are closely related by sequence identity, and more distantly related. Nucleic acids so identified can be translated into polypeptides and the polypeptides can be tested for activity. Identification of related sequences can also be achieved using polymerase chain reaction (PCR) and other amplification techniques suitable for cloning related nucleic acid sequences. Preferably PCR primers are selected to amplify portions of a nucleic acid sequence believed to be conserved (e.g., a catalytic domain, a DNA-binding domain, etc.). Again, nucleic acids are preferably amplified from a tissue-specific library (e.g., such as testis to find sequences related to OY-MC-4). One also can use expression cloning utilizing the antisera described herein to identify nucleic acids which encode related antigenic proteins in humans or other species using the SEREX procedure.

The term "stringent conditions" as used herein refers to parameters with which the art is familiar. Nucleic acid hybridization parameters may be found in references which compile such methods, e.g. *Molecular Cloning: A Laboratory Manual*, J. Sambrook, et al., eds., Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 1989, or *Current Protocols in Molecular Biology*, F.M. Ausubel, et al., eds., John Wiley & Sons, Inc., New York. More specifically, stringent conditions, as used herein, refers, for example, to hybridization at 65°C in hybridization buffer (3.5 x SSC, 0.02% Ficoll, 0.02% polyvinyl pyrrolidone, 0.02% Bovine Serum Albumin, 2.5mM NaH₂PO₄(pH7), 0.5% SDS, 2mM EDTA). SSC is 0.15M sodium chloride/0.15M sodium citrate, pH7; SDS is sodium dodecyl sulphate; and EDTA is ethylenediaminetetracetic acid. After hybridization, the membrane upon which the DNA is transferred is washed, for example, in 2 x SSC at room temperature and then at 0.1 - 0.5 x SSC/0.1 x SDS at temperatures up to 68°C.

There are other conditions, reagents, and so forth which can be used, which result in a

similar degree of stringency. The skilled artisan will be familiar with such conditions, and thus they are not given here. It will be understood, however, that the skilled artisan will be able to manipulate the conditions in a manner to permit the clear identification of homologs and alleles of cancer associated antigen nucleic acids of the invention (e.g., by using lower stringency conditions). The skilled artisan also is familiar with the methodology for screening cells and libraries for expression of such molecules which then are routinely isolated, followed by isolation of the pertinent nucleic acid molecule and sequencing.

In general homologs and alleles typically will share at least 75% nucleotide identity and/or at least 90% amino acid identity to the sequences of cancer associated antigen nucleic acid and polypeptides, respectively, in some instances will share at least 90% nucleotide identity and/or at least 95% amino acid identity and in still other instances will share at least 95% nucleotide identity and/or at least 99% amino acid identity. The homology can be calculated using various, publicly available software tools developed by NCBI (Bethesda, Maryland) that can be obtained through the internet (<ftp://ncbi.nlm.nih.gov/pub/>). Exemplary tools include the BLAST system available at <http://www.ncbi.nlm.nih.gov>, using default settings. Pairwise and ClustalW alignments (BLOSUM30 matrix setting) as well as Kyte-Doolittle hydropathic analysis can be obtained using the MacVector sequence analysis software (Oxford Molecular Group). Watson-Crick complements of the foregoing nucleic acids also are embraced by the invention.

In screening for cancer associated antigen genes, a Southern blot may be performed using the foregoing conditions, together with a radioactive probe. After washing the membrane to which the DNA is finally transferred, the membrane can be placed against X-ray film to detect the radioactive signal. In screening for the expression of cancer associated antigen nucleic acids, Northern blot hybridizations using the foregoing conditions (see also the Examples) can be performed on samples taken from cancer patients or subjects suspected of having a condition characterized by expression of cancer associated antigen genes. Amplification protocols such as polymerase chain reaction using primers which hybridize to the sequences presented also can be used for detection of the cancer associated antigen genes or expression thereof.

As used herein with respect to nucleic acids, the term "isolated" means: (i) amplified *in vitro* by, for example, polymerase chain reaction (PCR); (ii) recombinantly produced by cloning; (iii) purified, as by cleavage and gel separation; or (iv) synthesized by, for example, chemical synthesis. An isolated nucleic acid is one which is readily manipulable by

recombinant DNA techniques well known in the art. Thus, a nucleotide sequence contained in a vector in which 5' and 3' restriction sites are known or for which polymerase chain reaction (PCR) primer sequences have been disclosed is considered isolated but a nucleic acid sequence existing in its native state in its natural host is not. An isolated nucleic acid may be substantially purified, but need not be. For example, a nucleic acid that is isolated within a cloning or expression vector is not pure in that it may comprise only a tiny percentage of the material in the cell in which it resides. Such a nucleic acid is isolated, however, as the term is used herein because it is readily manipulable by standard techniques known to those of ordinary skill in the art. An isolated nucleic acid as used herein is not a naturally occurring chromosome.

As used herein with respect to polypeptides, "isolated" means separated from its native environment and present in sufficient quantity to permit its identification or use. Isolated, when referring to a protein or polypeptide, means, for example: (i) selectively produced by expression cloning or (ii) purified as by chromatography or electrophoresis. Isolated proteins or polypeptides may, but need not be, substantially pure. The term "substantially pure" means that the proteins or polypeptides are essentially free of other substances with which they may be found in nature or *in vivo* systems to an extent practical and appropriate for their intended use. Substantially pure polypeptides may be produced by techniques well known in the art. Because an isolated protein may be admixed with a pharmaceutically acceptable carrier in a pharmaceutical preparation, the protein may comprise only a small percentage by weight of the preparation. The protein is nonetheless isolated in that it has been separated from the substances with which it may be associated in living systems, i.e. isolated from other proteins.

The cancer associated genes correspond to SEQ ID Nos: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19 and 23. The preferred cancer associated antigens for the methods of diagnosis disclosed herein are those which are found to react with allogeneic cancer antisera (i.e. NA Group 5). Encoded polypeptides (e.g., proteins), peptides and antisera thereto are also preferred for diagnosis. Most preferably, the nucleic acid is SEQ ID NO:23 and the polypeptide is SEQ ID NO:24.

The invention also includes degenerate nucleic acids which include alternative codons to those present in the native materials. For example, serine residues are encoded by the codons TCA, AGT, TCC, TCG, TCT and AGC. Each of the six codons is equivalent for the purposes of encoding a serine residue. Thus, it will be apparent to one of ordinary skill in the art that any of the serine-encoding nucleotide triplets may be employed to direct the protein

synthesis apparatus, *in vitro* or *in vivo*, to incorporate a serine residue into an elongating cancer associated antigen polypeptide. Similarly, nucleotide sequence triplets which encode other amino acid residues include, but are not limited to: CCA, CCC, CCG and CCT (proline codons); CGA, CGC, CGG, CGT, AGA and AGG (arginine codons); ACA, ACC, ACG and
5 ACT (threonine codons); AAC and AAT (asparagine codons); and ATA, ATC and ATT (isoleucine codons). Other amino acid residues may be encoded similarly by multiple nucleotide sequences. Thus, the invention embraces degenerate nucleic acids that differ from the biologically isolated nucleic acids in codon sequence due to the degeneracy of the genetic code.

10 The invention also provides modified nucleic acid molecules which include additions, substitutions and deletions of one or more nucleotides. In preferred embodiments, these modified nucleic acid molecules and/or the polypeptides they encode retain at least one activity or function of the unmodified nucleic acid molecule and/or the polypeptides, such as antigenicity, enzymatic activity, receptor binding, formation of complexes by binding of
15 peptides by MHC class I and class II molecules, etc. In certain embodiments, the modified nucleic acid molecules encode modified polypeptides, preferably polypeptides having conservative amino acid substitutions as are described elsewhere herein. The modified nucleic acid molecules are structurally related to the unmodified nucleic acid molecules and in preferred embodiments are sufficiently structurally related to the unmodified nucleic acid
20 molecules so that the modified and unmodified nucleic acid molecules hybridize under stringent conditions known to one of skill in the art.

For example, modified nucleic acid molecules which encode polypeptides having single amino acid changes can be prepared. Each of these nucleic acid molecules can have one, two or three nucleotide substitutions exclusive of nucleotide changes corresponding to
25 the degeneracy of the genetic code as described herein. Likewise, modified nucleic acid molecules which encode polypeptides having two amino acid changes can be prepared which have, e.g., 2-6 nucleotide changes. Numerous modified nucleic acid molecules like these will be readily envisioned by one of skill in the art, including for example, substitutions of nucleotides in codons encoding amino acids 2 and 3, 2 and 4, 2 and 5, 2 and 6, and so on. In
30 the foregoing example, each combination of two amino acids is included in the set of modified nucleic acid molecules, as well as all nucleotide substitutions which code for the amino acid substitutions. Additional nucleic acid molecules that encode polypeptides having additional substitutions (i.e., 3 or more), additions or deletions (e.g., by introduction of a stop

codon or a splice site(s)) also can be prepared and are embraced by the invention as readily envisioned by one of ordinary skill in the art. Any of the foregoing nucleic acids or polypeptides can be tested by routine experimentation for retention of structural relation or activity to the nucleic acids and/or polypeptides disclosed herein.

5 The invention also provides isolated unique fragments of cancer associated antigen nucleic acid sequences or complements thereof. A unique fragment is one that is a 'signature' for the larger nucleic acid. It, for example, is long enough to assure that its precise sequence is not found in molecules within the human genome outside of the cancer associated antigen nucleic acids defined above (and human alleles). Those of ordinary skill in the art may apply
10 no more than routine procedures to determine if a fragment is unique within the human genome. Unique fragments, however, exclude fragments completely composed of the nucleotide sequences of any of GenBank accession numbers listed in Table 8 or other previously published sequences as of the filing date of the priority documents for sequences listed in a respective priority document or the filing date of this application for sequences
15 listed for the first time in this application which overlap the sequences of the invention.

A fragment which is completely composed of the sequence described in the foregoing GenBank deposits is one which does not include any of the nucleotides unique to the sequences of the invention. Thus, a unique fragment must contain a nucleotide sequence other than the exact sequence of those in GenBank or fragments thereof. The difference may
20 be an addition, deletion or substitution with respect to the GenBank sequence or it may be a sequence wholly separate from the GenBank sequence.

Unique fragments can be used as probes in Southern and Northern blot assays to identify such nucleic acids, or can be used in amplification assays such as those employing PCR. As known to those skilled in the art, large probes such as 200, 250, 300 or more
25 nucleotides are preferred for certain uses such as Southern and Northern blots, while smaller fragments will be preferred for uses such as PCR. Unique fragments also can be used to produce fusion proteins for generating antibodies or determining binding of the polypeptide fragments, or for generating immunoassay components. Likewise, unique fragments can be employed to produce nonfused fragments of the cancer associated antigen polypeptides,
30 useful, for example, in the preparation of antibodies, and in immunoassays. Unique fragments further can be used as antisense molecules to inhibit the expression of cancer associated antigen nucleic acids and polypeptides, particularly for therapeutic purposes as described in greater detail below.

As will be recognized by those skilled in the art, the size of the unique fragment will depend upon its conservancy in the genetic code. Thus, some regions of cancer associated antigen sequences and complements thereof will require longer segments to be unique while others will require only short segments, typically between 12 and 32 nucleotides (e.g. 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31 and 32 or more bases long, up to the entire length of the disclosed sequence. As mentioned above, this disclosure intends to embrace each and every fragment of each sequence, beginning at the first nucleotide, the second nucleotide and so on, up to 8 nucleotides short of the end, and ending anywhere from nucleotide number 8, 9, 10 and so on for each sequence, up to the very last nucleotide (provided the sequence is unique as described above).

Many segments of the polypeptide coding region of novel cancer associated antigen nucleic acids, or complements thereof, that is 18 or more nucleotides in length will be unique. Those skilled in the art are well versed in methods for selecting such sequences, typically on the basis of the ability of the unique fragment to selectively distinguish the sequence of interest from other sequences in the human genome of the fragment to those on known databases typically is all that is necessary, although *in vitro* confirmatory hybridization and sequencing analysis may be performed.

Especially preferred include nucleic acids encoding a series of epitopes, known as "polytopes". The epitopes can be arranged in sequential or overlapping fashion (*see, e.g.,* Thomson et al., *Proc. Natl. Acad. Sci. USA* 92:5845-5849, 1995; Gilbert et al., *Nature Biotechnol.* 15:1280-1284, 1997), with or without the natural flanking sequences, and can be separated by unrelated linker sequences if desired. The polytope is processed to generated individual epitopes which are recognized by the immune system for generation of immune responses.

Thus, for example, peptides derived from a polypeptide having an amino acid sequence encoded by one of the nucleic acid disclosed herein, and which are presented by MHC molecules and recognized by CTL or T helper lymphocytes, can be combined with peptides from one or more other cancer associated antigens (e.g. by preparation of hybrid nucleic acids or polypeptides) to form "polytopes". The two or more peptides (or nucleic acids encoding the peptides) can be selected from those described herein, or they can include one or more peptides of previously known cancer associated antigens. Exemplary cancer associated peptide antigens that can be administered to induce or enhance an immune response are derived from tumor associated genes and encoded proteins including MAGE-A1,

MAGE-A2, MAGE-A3, MAGE-A4, MAGE-A5, MAGE-A6, MAGE-A7, MAGE-A8, MAGE-A9, MAGE-A10, MAGE-A11, MAGE-A12, MAGE-A13, GAGE-1, GAGE-2, GAGE-3, GAGE-4, GAGE-5, GAGE-6, GAGE-7, GAGE-8, BAGE-1, RAGE-1, LB33/MUM-1, PRAME, NAG, MAGE-B2, MAGE-B3, MAGE-B4, tyrosinase, brain
5 glycogen phosphorylase, Melan-A, MAGE-C1, MAGE-C2, NY-ESO-1, LAGE-1, SSX-1, SSX-2 (HOM-MEL-40), SSX-4, SSX-5, SCP-1 and CT-7. See, for example, PCT application publication no. WO96/10577. Other examples will be known to one of ordinary skill in the art (for example, see Coulie, *Stem Cells* 13:393-403, 1995), and can be used in the invention in a like manner as those disclosed herein. One of ordinary skill in the art can prepare
10 polypeptides comprising one or more peptides and one or more of the foregoing cancer associated peptides, or nucleic acids encoding such polypeptides, according to standard procedures of molecular biology.

Thus polytopes are groups of two or more potentially immunogenic or immune response stimulating peptides which can be joined together in various arrangements (e.g.
15 concatenated, overlapping). The polytope (or nucleic acid encoding the polytope) can be administered in a standard immunization protocol, e.g. to animals, to test the effectiveness of the polytope in stimulating, enhancing and/or provoking an immune response.

The peptides can be joined together directly or via the use of flanking sequences to form polytopes, and the use of polytopes as vaccines is well known in the art (see, e.g.,
20 Thomson et al., *Proc. Acad. Natl. Acad. Sci USA* 92(13):5845-5849, 1995; Gilbert et al., *Nature Biotechnol.* 15(12):1280-1284, 1997; Thomson et al., *J. Immunol.* 157(2):822-826, 1996; Tam et al., *J. Exp. Med.* 171(1):299-306, 1990). For example, Tam showed that polytopes consisting of both MHC class I and class II binding epitopes successfully generated antibody and protective immunity in a mouse model. Tam also demonstrated that polytopes
25 comprising "strings" of epitopes are processed to yield individual epitopes which are presented by MHC molecules and recognized by CTLs. Thus polytopes containing various numbers and combinations of epitopes can be prepared and tested for recognition by CTLs and for efficacy in increasing an immune response.

It is known that tumors express a set of tumor antigens, of which only certain subsets
30 may be expressed in the tumor of any given patient. Polytopes can be prepared which correspond to the different combination of epitopes representing the subset of tumor rejection antigens expressed in a particular patient. Polytopes also can be prepared to reflect a broader spectrum of tumor rejection antigens known to be expressed by a tumor type. Polytopes can

be introduced to a patient in need of such treatment as polypeptide structures, or via the use of nucleic acid delivery systems known in the art (see, e.g., Allsopp et al., *Eur. J. Immunol.* 26(8):1951-1959, 1996). Adenovirus, pox virus, Ty-virus like particles, adeno-associated virus, plasmids, bacteria, etc. can be used in such delivery. One can test the polytope delivery systems in mouse models to determine efficacy of the delivery system. The systems also can be tested in human clinical trials.

In instances in which a MHC class I molecule presents tumor rejection antigens derived from cancer associated nucleic acids, the expression vector may also include a nucleic acid sequence coding for the MHC molecule that presents any particular tumor rejection antigen derived from these nucleic acids and polypeptides. Alternatively, the nucleic acid sequence coding for such a MHC molecule can be contained within a separate expression vector. In a situation where the vector contains both coding sequences, the single vector can be used to transfect a cell which does not normally express either one. Where the coding sequences for a cancer associated antigen precursor and the MHC molecule which presents it are contained on separate expression vectors, the expression vectors can be cotransfected. The cancer associated antigen precursor coding sequence may be used alone, when, e.g. the host cell already expresses a MHC molecule which presents a cancer associated antigen derived from precursor molecules. Of course, there is no limit on the particular host cell which can be used. As the vectors which contain the two coding sequences may be used in any antigen-presenting cells if desired, and the gene for cancer associated antigen precursor can be used in host cells which do not express a MHC molecule which presents a cancer associated antigen. Further, cell-free transcription systems may be used in lieu of cells.

As mentioned above, the invention embraces antisense oligonucleotides that selectively bind to a nucleic acid molecule encoding a cancer associated antigen polypeptide, to reduce the expression of cancer associated antigens. This is desirable in virtually any medical condition wherein a reduction of expression of cancer associated antigens is desirable, e.g., in the treatment of cancer. This is also useful for *in vitro* or *in vivo* testing of the effects of a reduction of expression of one or more cancer associated antigens.

As used herein, the term "antisense oligonucleotide" or "antisense" describes an oligonucleotide that is an oligoribonucleotide, oligodeoxyribonucleotide, modified oligoribonucleotide, or modified oligodeoxyribonucleotide which hybridizes under physiological conditions to DNA comprising a particular gene or to an mRNA transcript of that gene and, thereby, inhibits the transcription of that gene and/or the translation of that

mRNA. The antisense molecules are designed so as to interfere with transcription or translation of a target gene upon hybridization with the target gene or transcript. Those skilled in the art will recognize that the exact length of the antisense oligonucleotide and its degree of complementarity with its target will depend upon the specific target selected, including the sequence of the target and the particular bases which comprise that sequence. It is preferred that the antisense oligonucleotide be constructed and arranged so as to bind selectively with the target under physiological conditions, i.e., to hybridize substantially more to the target sequence than to any other sequence in the target cell under physiological conditions. Based upon the sequences of nucleic acids encoding cancer associated antigens, or upon allelic or homologous genomic and/or cDNA sequences, one of skill in the art can easily choose and synthesize any of a number of appropriate antisense molecules for use in accordance with the present invention. In order to be sufficiently selective and potent for inhibition, such antisense oligonucleotides should comprise at least 10 and, more preferably, at least 15 consecutive bases which are complementary to the target, although in certain cases modified oligonucleotides as short as 7 bases in length have been used successfully as antisense oligonucleotides (Wagner et al., *Nature Biotechnol.* 14:840-844, 1996). Most preferably, the antisense oligonucleotides comprise a complementary sequence of 20-30 bases. Although oligonucleotides may be chosen which are antisense to any region of the gene or mRNA transcripts, in preferred embodiments the antisense oligonucleotides correspond to N-terminal or 5' upstream sites such as translation initiation, transcription initiation or promoter sites. In addition, 3'-untranslated regions may be targeted. Targeting to mRNA splicing sites has also been used in the art but may be less preferred if alternative mRNA splicing occurs. In addition, the antisense is targeted, preferably, to sites in which mRNA secondary structure is not expected (see, e.g., Sainio et al., *Cell Mol. Neurobiol.* 14(5):439-457, 1994) and at which proteins are not expected to bind. Finally, although the listed sequences are cDNA sequences, one of ordinary skill in the art may easily derive the genomic DNA corresponding to the cDNA of a cancer associated antigen. Thus, the present invention also provides for antisense oligonucleotides which are complementary to the genomic DNA corresponding to nucleic acids encoding cancer associated antigens. Similarly, antisense to allelic cDNAs, homologous cDNAs (e.g., human) and genomic DNAs are enabled without undue experimentation.

In one set of embodiments, the antisense oligonucleotides of the invention may be composed of "natural" deoxyribonucleotides, ribonucleotides, or any combination thereof.

That is, the 5' end of one native nucleotide and the 3' end of another native nucleotide may be covalently linked, as in natural systems, via a phosphodiester internucleoside linkage. These oligonucleotides may be prepared by art recognized methods which may be carried out manually or by an automated synthesizer. They also may be produced recombinantly by
5 vectors.

In preferred embodiments, however, the antisense oligonucleotides of the invention also may include "modified" oligonucleotides. That is, the oligonucleotides may be modified in a number of ways which do not prevent them from hybridizing to their target but which enhance their stability or targeting or which otherwise enhance their therapeutic effectiveness.

10 The term "modified oligonucleotide" as used herein describes an oligonucleotide in which (1) at least two of its nucleotides are covalently linked via a synthetic internucleoside linkage (i.e., a linkage other than a phosphodiester linkage between the 5' end of one nucleotide and the 3' end of another nucleotide) and/or (2) a chemical group not normally associated with nucleic acids has been covalently attached to the oligonucleotide. Preferred
15 synthetic internucleoside linkages are phosphorothioates, alkylphosphonates, phosphorodithioates, phosphate esters, alkylphosphonothioates, phosphoramidates, carbamates, carbonates, phosphate triesters, acetamides, carboxymethyl esters and peptides.

The term "modified oligonucleotide" also encompasses oligonucleotides with a covalently modified base and/or sugar. For example, modified oligonucleotides include
20 oligonucleotides having backbone sugars which are covalently attached to low molecular weight organic groups other than a hydroxyl group at the 3' position and other than a phosphate group at the 5' position. Thus modified oligonucleotides may include a 2'-O-alkylated ribose group. In addition, modified oligonucleotides may include sugars such as arabinose instead of ribose. The present invention, thus, contemplates pharmaceutical
25 preparations containing modified antisense molecules that are complementary to and hybridizable with, under physiological conditions, nucleic acids encoding cancer associated antigen polypeptides, together with pharmaceutically acceptable carriers.

As used herein, a "vector" may be any of a number of nucleic acids into which a desired sequence may be inserted by restriction and ligation for transport between different
30 genetic environments or for expression in a host cell. Vectors are typically composed of DNA although RNA vectors are also available. Vectors include, but are not limited to, plasmids, phagemids and virus genomes. A cloning vector is one which is able to replicate autonomously or integrated in the genome in a host cell, and which is further characterized by

one or more endonuclease restriction sites at which the vector may be cut in a determinable fashion and into which a desired DNA sequence may be ligated such that the new recombinant vector retains its ability to replicate in the host cell. In the case of plasmids, replication of the desired sequence may occur many times as the plasmid increases in copy number within the host bacterium or just a single time per host before the host reproduces by mitosis. In the case of phage, replication may occur actively during a lytic phase or passively during a lysogenic phase. An expression vector is one into which a desired DNA sequence may be inserted by restriction and ligation such that it is operably joined to regulatory sequences and may be expressed as an RNA transcript. Vectors may further contain one or more marker sequences suitable for use in the identification of cells which have or have not been transformed or transfected with the vector. Markers include, for example, genes encoding proteins which increase or decrease either resistance or sensitivity to antibiotics or other compounds, genes which encode enzymes whose activities are detectable by standard assays known in the art (e.g., β -galactosidase, luciferase or alkaline phosphatase), and genes which visibly affect the phenotype of transformed or transfected cells, hosts, colonies or plaques (e.g., green fluorescent protein). Preferred vectors are those capable of autonomous replication and expression of the structural gene products present in the DNA segments to which they are operably joined.

As used herein, a coding sequence and regulatory sequences are said to be “operably” joined when they are covalently linked in such a way as to place the expression or transcription of the coding sequence under the influence or control of the regulatory sequences. If it is desired that the coding sequences be translated into a functional protein, two DNA sequences are said to be operably joined if induction of a promoter in the 5’ regulatory sequences results in the transcription of the coding sequence and if the nature of the linkage between the two DNA sequences does not (1) result in the introduction of a frame-shift mutation, (2) interfere with the ability of the promoter region to direct the transcription of the coding sequences, or (3) interfere with the ability of the corresponding RNA transcript to be translated into a protein. Thus, a promoter region would be operably joined to a coding sequence if the promoter region were capable of effecting transcription of that DNA sequence such that the resulting transcript might be translated into the desired protein or polypeptide.

The precise nature of the regulatory sequences needed for gene expression may vary between species or cell types, but shall in general include, as necessary, 5’ non-transcribed and 5’ non-translated sequences involved with the initiation of transcription and translation

respectively, such as a TATA box, capping sequence, CAAT sequence, and the like.

Especially, such 5' non-transcribed regulatory sequences will include a promoter region which includes a promoter sequence for transcriptional control of the operably joined gene.

Regulatory sequences may also include enhancer sequences or upstream activator sequences

5 as desired. The vectors of the invention may optionally include 5' leader or signal sequences. The choice and design of an appropriate vector is within the ability and discretion of one of ordinary skill in the art.

Expression vectors containing all the necessary elements for expression are commercially available and known to those skilled in the art. See, e.g., Sambrook et al.,
10 *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press, 1989. Cells are genetically engineered by the introduction into the cells of heterologous DNA (RNA) encoding a cancer associated antigen polypeptide or fragment or variant thereof. That heterologous DNA (RNA) is placed under operable control of transcriptional elements to permit the expression of the heterologous DNA in the host cell.

15 Preferred systems for mRNA expression in mammalian cells are those such as pcDNA3.1 or pRc/CMV (available from Invitrogen, Carlsbad, CA) that contain a selectable marker such as a gene that confers G418 resistance (which facilitates the selection of stably transfected cell lines) and the human cytomegalovirus (CMV) enhancer-promoter sequences. Additionally, suitable for expression in primate or canine cell lines is the pCEP4 vector
20 (Invitrogen), which contains an Epstein Barr Virus (EBV) origin of replication, facilitating the maintenance of plasmid as a multicopy extrachromosomal element. Another expression vector is the pEF-BOS plasmid containing the promoter of polypeptide Elongation Factor 1 α , which stimulates efficiently transcription *in vitro*. The plasmid is described by Mishizuma and Nagata (*Nuc. Acids Res.* 18:5322, 1990), and its use in transfection experiments is
25 disclosed by, for example, Demoulin (*Mol. Cell. Biol.* 16:4710-4716, 1996). Still another preferred expression vector is an adenovirus, described by Stratford-Perricaudet, which is defective for E1 and E3 proteins (*J. Clin. Invest.* 90:626-630, 1992). The use of the adenovirus as an Adeno.P1A recombinant for the expression of an antigen is disclosed by Warnier et al., in intradermal injection in mice for immunization against P1A (*Int. J. Cancer*,
30 67:303-310, 1996). Additional vectors for delivery of nucleic acid are provided below.

The invention also embraces so-called expression kits, which allow the artisan to prepare a desired expression vector or vectors. Such expression kits include at least separate portions of a vector and one or more of the previously discussed cancer associated antigen

nucleic acid molecules. Other components may be added, as desired, as long as the previously mentioned nucleic acid molecules, which are required, are included. The invention also includes kits for amplification of a cancer associated antigen nucleic acid, including at least one pair of amplification primers which hybridize to a cancer associated antigen nucleic acid. The primers preferably are 12-32 nucleotides in length and are non-overlapping to prevent formation of "primer-dimers". One of the primers will hybridize to one strand of the cancer associated antigen nucleic acid and the second primer will hybridize to the complementary strand of the cancer associated antigen nucleic acid, in an arrangement which permits amplification of the cancer associated antigen nucleic acid. Selection of appropriate primer pairs is standard in the art. For example, the selection can be made with assistance of a computer program designed for such a purpose, optionally followed by testing the primers for amplification specificity and efficiency.

The invention also permits the construction of cancer associated antigen gene "knock-outs" and transgenics in cells and in animals, providing materials for studying certain aspects of cancer and immune system responses to cancer.

The invention also provides isolated polypeptides (including whole proteins and partial proteins) encoded by the foregoing cancer associated antigen nucleic acids. Such polypeptides are useful, for example, alone or as fusion proteins to generate antibodies, as components of an immunoassay or diagnostic assay or as therapeutics. Cancer associated antigen polypeptides can be isolated from biological samples including tissue or cell homogenates, and can also be expressed recombinantly in a variety of prokaryotic and eukaryotic expression systems by constructing an expression vector appropriate to the expression system, introducing the expression vector into the expression system, and isolating the recombinantly expressed protein. Short polypeptides, including antigenic peptides (such as are presented by MHC molecules on the surface of a cell for immune recognition) also can be synthesized chemically using well-established methods of peptide synthesis.

A unique fragment of a cancer associated antigen polypeptide, in general, has the features and characteristics of unique fragments as discussed above in connection with nucleic acids. As will be recognized by those skilled in the art, the size of the unique fragment will depend upon factors such as whether the fragment constitutes a portion of a conserved protein domain. Thus, some regions of cancer associated antigens will require longer segments to be unique while others will require only short segments, typically between 5 and 12 amino acids (e.g. 5, 6, 7, 8, 9, 10, 11 or 12 or more amino acids including each integer up to the full

length).

Human homologs of cancer associated antigen polypeptides are related in sequence to the cancer associated antigens described herein, and preferably are also related in function. Preferably the homologs are 90% or more identical to one or more portions of the amino acid sequence of the cancer associated antigens, more preferably are 95% or more identical, and still more preferably are 99% or more identical. Most preferably, the homologs contains at least one fragment of 10 or more amino acids that are identical to the corresponding amino acids of the cancer associated antigens. In some embodiments a human homolog has the same or similar activity or function as a cancer associated antigen. Activities and functions include, but are not limited to, enzymatic activity, recognition by antibodies, DNA binding activity, transcriptional activity, binding to MHC molecules, and the like.

Unique fragments and human homologs of a polypeptide preferably are those fragments and homologs which retain a distinct functional capability of the polypeptide. Functional capabilities which can be retained in a unique fragment of a polypeptide include interaction with antibodies, interaction with other polypeptides or fragments thereof, selective binding of nucleic acids or proteins, and enzymatic activity. One important activity is the ability to act as a signature for identifying the polypeptide. Another is the ability to complex with MHC and to provoke in a mammal, preferably a human, an immune response. Those skilled in the art are well versed in methods for selecting unique amino acid sequences, typically on the basis of the ability of the unique fragment to selectively distinguish the sequence of interest from non-family members. A comparison of the sequence of the fragment to those on known databases typically is all that is necessary.

The invention embraces variants of the cancer associated antigen polypeptides described above. As used herein, a "variant" of a cancer associated antigen polypeptide is a polypeptide which contains one or more modifications to the primary amino acid sequence of a cancer associated antigen polypeptide. Modifications which create a cancer associated antigen variant can be made to a cancer associated antigen polypeptide 1) to reduce or eliminate an activity of a cancer associated antigen polypeptide; 2) to enhance a property of a cancer associated antigen polypeptide, such as protein stability in an expression system or the stability of protein-protein binding; 3) to provide a novel activity or property to a cancer associated antigen polypeptide, such as addition of an antigenic epitope or addition of a detectable moiety; or 4) to provide equivalent or better binding to a MHC molecule. Modifications to a cancer associated antigen polypeptide are typically made to the nucleic

acid which encodes the cancer associated antigen polypeptide, and can include deletions, point mutations, truncations, amino acid substitutions and additions of amino acids or non-amino acid moieties. Alternatively, modifications can be made directly to the polypeptide, such as by cleavage, addition of a linker molecule, addition of a detectable moiety, such as biotin, addition of a fatty acid, and the like. Modifications also embrace fusion proteins comprising all or part of the cancer associated antigen amino acid sequence. One of skill in the art will be familiar with methods for predicting the effect on protein conformation of a change in protein sequence, and can thus “design” a variant cancer associated antigen polypeptide according to known methods. One example of such a method is described by Dahiyat and Mayo in *Science* 278:82-87, 1997, whereby proteins can be designed *de novo*. The method can be applied to a known protein to vary only a portion of the polypeptide sequence. By applying the computational methods of Dahiyat and Mayo, specific variants of a cancer associated antigen polypeptide can be proposed and tested to determine whether the variant retains a desired conformation.

In general, variants include cancer associated antigen polypeptides which are modified specifically to alter a feature of the polypeptide unrelated to its desired physiological activity. For example, cysteine residues can be substituted or deleted to prevent unwanted disulfide linkages. Similarly, certain amino acids can be changed to enhance expression of a cancer associated antigen polypeptide by eliminating proteolysis by proteases in an expression system (e.g., dibasic amino acid residues in yeast expression systems in which KEX2 protease activity is present).

Mutations of a nucleic acid which encode a cancer associated antigen polypeptide preferably preserve the amino acid reading frame of the coding sequence, and preferably do not create regions in the nucleic acid which are likely to hybridize to form secondary structures, such as hairpins or loops, which can be deleterious to expression of the variant polypeptide.

Mutations can be made by selecting an amino acid substitution, or by random mutagenesis of a selected site in a nucleic acid which encodes the polypeptide. Variant polypeptides are then expressed and tested for one or more activities to determine which mutation provides a variant polypeptide with the desired properties. Further mutations can be made to variants (or to non-variant cancer associated antigen polypeptides) which are silent as to the amino acid sequence of the polypeptide, but which provide preferred codons for translation in a particular host. The preferred codons for translation of a nucleic acid in, e.g.,

E. coli, are well known to those of ordinary skill in the art. Still other mutations can be made to the noncoding sequences of a cancer associated antigen gene or cDNA clone to enhance expression of the polypeptide. The activity of variants of cancer associated antigen polypeptides can be tested by cloning the gene encoding the variant cancer associated antigen polypeptide into a bacterial or mammalian expression vector, introducing the vector into an appropriate host cell, expressing the variant cancer associated antigen polypeptide, and testing for a functional capability of the cancer associated antigen polypeptides as disclosed herein. For example, the variant cancer associated antigen polypeptide can be tested for reaction with autologous or allogeneic sera as described in the Examples. Preparation of other variant polypeptides may favor testing of other activities, as will be known to one of ordinary skill in the art.

The skilled artisan will also realize that conservative amino acid substitutions may be made in cancer associated antigen polypeptides to provide functionally equivalent variants of the foregoing polypeptides, i.e., the variants retain the functional capabilities of the cancer associated antigen polypeptides. As used herein, a "conservative amino acid substitution" refers to an amino acid substitution which does not alter the relative charge or size characteristics of the protein in which the amino acid substitution is made. Variants can be prepared according to methods for altering polypeptide sequence known to one of ordinary skill in the art such as are found in references which compile such methods, e.g. *Molecular Cloning: A Laboratory Manual*, J. Sambrook, et al., eds., Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 1989, or *Current Protocols in Molecular Biology*, F.M. Ausubel, et al., eds., John Wiley & Sons, Inc., New York. Exemplary functionally equivalent variants of the cancer associated antigen polypeptides include conservative amino acid substitutions of in the amino acid sequences of proteins disclosed herein. Conservative substitutions of amino acids include substitutions made amongst amino acids within the following groups: (a) M, I, L, V; (b) F, Y, W; (c) K, R, H; (d) A, G; (e) S, T; (f) Q, N; and (g) E, D.

For example, upon determining that a peptide derived from a cancer associated antigen polypeptide is presented by an MHC molecule and recognized by antibodies or CTLs, one can make conservative amino acid substitutions to the amino acid sequence of the peptide, particularly at residues which are thought not to be direct contact points with the MHC molecule. For example, methods for identifying functional variants of HLA class II binding peptides are provided in a published PCT application of Strominger and Wucherpennig

(PCT/US96/03182). Peptides bearing one or more amino acid substitutions also can be tested for concordance with known HLA/MHC motifs prior to synthesis using, e.g. the computer program described by D'Amaro and Drijfhout (D'Amaro et al., *Human Immunol.* 43:13-18, 1995; Drijfhout et al., *Human Immunol.* 43:1-12, 1995). The substituted peptides can then be tested for binding to the MHC molecule and recognition by antibodies or CTLs when bound to MHC. These variants can be tested for improved stability and are useful, *inter alia*, in vaccine compositions.

Conservative amino-acid substitutions in the amino acid sequence of cancer associated antigen polypeptides to produce functionally equivalent variants of cancer associated antigen polypeptides typically are made by alteration of a nucleic acid encoding a cancer associated antigen polypeptide. Such substitutions can be made by a variety of methods known to one of ordinary skill in the art. For example, amino acid substitutions may be made by PCR-directed mutation, site-directed mutagenesis according to the method of Kunkel (Kunkel, *Proc. Nat. Acad. Sci. U.S.A.* 82: 488-492, 1985), or by chemical synthesis of a gene encoding a cancer associated antigen polypeptide. Where amino acid substitutions are made to a small unique fragment of a cancer associated antigen polypeptide, such as an antigenic epitope recognized by autologous or allogeneic sera or cytolytic T lymphocytes, the substitutions can be made by directly synthesizing the peptide. The activity of functionally equivalent fragments of cancer associated antigen polypeptides can be tested by cloning the gene encoding the altered cancer associated antigen polypeptide into a bacterial or mammalian expression vector, introducing the vector into an appropriate host cell, expressing the altered cancer associated antigen polypeptide, and testing for a functional capability of the cancer associated antigen polypeptides as disclosed herein. Peptides which are chemically synthesized can be tested directly for function, e.g., for binding to antisera recognizing associated antigens.

The invention as described herein has a number of uses, some of which are described elsewhere herein. First, the invention permits isolation of the cancer associated antigen protein molecules. A variety of methodologies well-known to the skilled practitioner can be utilized to obtain isolated cancer associated antigen molecules. The polypeptide may be purified from cells which naturally produce the polypeptide by chromatographic means or immunological recognition. Alternatively, an expression vector may be introduced into cells to cause production of the polypeptide. In another method, mRNA transcripts may be microinjected or otherwise introduced into cells to cause production of the encoded polypeptide. Translation of mRNA in cell-free extracts such as the reticulocyte lysate system

also may be used to produce polypeptide. Those skilled in the art also can readily follow known methods for isolating cancer associated antigen polypeptides. These include, but are not limited to, immunochromatography, HPLC, size-exclusion chromatography, ion-exchange chromatography and immune-affinity chromatography.

5 Identification of the cancer associated antigens also permits the identification and isolation of related human sequences (e.g. human homologs such as SEQ ID NOs:23 and 24), including immunogenic fragments of polypeptides and nucleic acids which encode such fragments. The isolation of homologs of different species is described elsewhere herein, and exemplified in Example 8.

10 The isolation and identification of cancer associated antigen genes also makes it possible for the artisan to diagnose a disorder characterized by expression of cancer associated antigens. These methods involve determining expression of one or more cancer associated antigen nucleic acids, and/or encoded cancer associated antigen polypeptides and/or peptides derived therefrom. In the former situation, such determinations can be carried out via any
15 standard nucleic acid determination assay, including the polymerase chain reaction, or assaying with labeled hybridization probes. In the latter situation, such determinations can be carried out by screening patient antisera for recognition of the polypeptide.

The invention also makes it possible isolate proteins which bind to cancer associated antigens as disclosed herein, including antibodies and cellular binding partners of the cancer
20 associated antigens. Additional uses are described further herein.

The invention also provides, in certain embodiments, "dominant negative" polypeptides derived from cancer associated antigen polypeptides. A dominant negative polypeptide is an inactive variant of a protein, which, by interacting with the cellular machinery, displaces an active protein from its interaction with the cellular machinery or
25 competes with the active protein, thereby reducing the effect of the active protein. For example, a dominant negative receptor which binds a ligand but does not transmit a signal in response to binding of the ligand can reduce the biological effect of expression of the ligand. Likewise, a dominant negative catalytically-inactive kinase which interacts normally with target proteins but does not phosphorylate the target proteins can reduce phosphorylation of
30 the target proteins in response to a cellular signal. Similarly, a dominant negative transcription factor which binds to a promoter site in the control region of a gene but does not increase gene transcription can reduce the effect of a normal transcription factor by occupying promoter binding sites without increasing transcription.

The end result of the expression of a dominant negative polypeptide in a cell is a reduction in function of active proteins. One of ordinary skill in the art can assess the potential for a dominant negative variant of a protein, and using standard mutagenesis techniques to create one or more dominant negative variant polypeptides. For example, given the teachings contained herein of cancer associated antigens, especially those which are similar to known proteins which have known activities (e.g., OY-MC-4 (*pem*)), one of ordinary skill in the art can modify the sequence of the cancer associated antigens by site-specific mutagenesis, scanning mutagenesis, partial gene deletion or truncation, and the like. See, e.g., U.S. Patent No. 5,580,723 and Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press, 1989. The skilled artisan then can test the population of mutagenized polypeptides for diminution in a selected and/or for retention of such an activity. Other similar methods for creating and testing dominant negative variants of a protein will be apparent to one of ordinary skill in the art.

The invention also involves agents such as polypeptides which bind to cancer associated antigen polypeptides. Such binding agents can be used, for example, in screening assays to detect the presence or absence of cancer associated antigen polypeptides and complexes of cancer associated antigen polypeptides and their binding partners and in purification protocols to isolated cancer associated antigen polypeptides and complexes of cancer associated antigen polypeptides and their binding partners. Such agents also can be used to inhibit the native activity of the cancer associated antigen polypeptides, for example, by binding to such polypeptides.

The invention, therefore, embraces peptide binding agents which, for example, can be antibodies or fragments of antibodies having the ability to selectively bind to cancer associated antigen polypeptides. Antibodies include polyclonal and monoclonal antibodies, prepared according to conventional methodology.

Significantly, as is well-known in the art, only a small portion of an antibody molecule, the paratope, is involved in the binding of the antibody to its epitope (see, in general, Clark, W.R. (1986) The Experimental Foundations of Modern Immunology Wiley & Sons, Inc., New York; Roitt, I. (1991) Essential Immunology, 7th Ed., Blackwell Scientific Publications, Oxford). The pFc' and Fc regions, for example, are effectors of the complement cascade but are not involved in antigen binding. An antibody from which the pFc' region has been enzymatically cleaved, or which has been produced without the pFc' region, designated an F(ab')₂ fragment, retains both of the antigen binding sites of an intact antibody. Similarly,

an antibody from which the Fc region has been enzymatically cleaved, or which has been produced without the Fc region, designated an Fab fragment, retains one of the antigen binding sites of an intact antibody molecule. Proceeding further, Fab fragments consist of a covalently bound antibody light chain and a portion of the antibody heavy chain denoted Fd.

5 The Fd fragments are the major determinant of antibody specificity (a single Fd fragment may be associated with up to ten different light chains without altering antibody specificity) and Fd fragments retain epitope-binding ability in isolation.

Within the antigen-binding portion of an antibody, as is well-known in the art, there are complementarity determining regions (CDRs), which directly interact with the epitope of the antigen, and framework regions (FRs), which maintain the tertiary structure of the paratope (see, in general, Clark, 1986; Roitt, 1991). In both the heavy chain Fd fragment and the light chain of IgG immunoglobulins, there are four framework regions (FR1 through FR4) separated respectively by three complementarity determining regions (CDR1 through CDR3).

The CDRs, and in particular the CDR3 regions, and more particularly the heavy chain CDR3, are largely responsible for antibody specificity.

It is now well-established in the art that the non-CDR regions of a mammalian antibody may be replaced with similar regions of conspecific or heterospecific antibodies while retaining the epitopic specificity of the original antibody. This is most clearly manifested in the development and use of "humanized" antibodies in which non-human CDRs are covalently joined to human FR and/or Fc/pFc' regions to produce a functional antibody. See, e.g., U.S. patents 4,816,567, 5,225,539, 5,585,089, 5,693,762 and 5,859,205.

Thus, for example, PCT International Publication Number WO 92/04381 teaches the production and use of humanized murine RSV antibodies in which at least a portion of the murine FR regions have been replaced by FR regions of human origin. Such antibodies, including fragments of intact antibodies with antigen-binding ability, are often referred to as "chimeric" antibodies.

Thus, as will be apparent to one of ordinary skill in the art, the present invention also provides for F(ab')₂, Fab, Fv and Fd fragments; chimeric antibodies in which the Fc and/or FR and/or CDR1 and/or CDR2 and/or light chain CDR3 regions have been replaced by homologous human or non-human sequences; chimeric F(ab')₂ fragment antibodies in which the FR and/or CDR1 and/or CDR2 and/or light chain CDR3 regions have been replaced by homologous human or non-human sequences; chimeric Fab fragment antibodies in which the FR and/or CDR1 and/or CDR2 and/or light chain CDR3 regions have been replaced by

homologous human or non-human sequences; and chimeric Fd fragment antibodies in which the FR and/or CDR1 and/or CDR2 regions have been replaced by homologous human or non-human sequences. The present invention also includes so-called single chain antibodies.

Thus, the invention involves polypeptides of numerous size and type that bind specifically to cancer associated antigen polypeptides, and complexes of both cancer associated antigen polypeptides and their binding partners. These polypeptides may be derived also from sources other than antibody technology. For example, such polypeptide binding agents can be provided by degenerate peptide libraries which can be readily prepared in solution, in immobilized form or as phage display libraries. Combinatorial libraries also can be synthesized of peptides containing one or more amino acids. Libraries further can be synthesized of peptoids and non-peptide synthetic moieties.

Phage display can be particularly effective in identifying binding peptides useful according to the invention. Briefly, one prepares a phage library (using e.g. m13, fd, or lambda phage), displaying inserts from 4 to about 80 amino acid residues using conventional procedures. The inserts may represent, for example, a completely degenerate or biased array. One then can select phage-bearing inserts which bind to the cancer associated antigen polypeptide. This process can be repeated through several cycles of reselection of phage that bind to the cancer associated antigen polypeptide. Repeated rounds lead to enrichment of phage bearing particular sequences. DNA sequence analysis can be conducted to identify the sequences of the expressed polypeptides. The minimal linear portion of the sequence that binds to the cancer associated antigen polypeptide can be determined. One can repeat the procedure using a biased library containing inserts containing part or all of the minimal linear portion plus one or more additional degenerate residues upstream or downstream thereof. Yeast two-hybrid screening methods also may be used to identify polypeptides that bind to the cancer associated antigen polypeptides. Thus, the cancer associated antigen polypeptides of the invention, or a fragment thereof, can be used to screen peptide libraries, including phage display libraries, to identify and select peptide binding partners of the cancer associated antigen polypeptides of the invention. Such molecules can be used, as described, for screening assays, for purification protocols, for interfering directly with the functioning of cancer associated antigen and for other purposes that will be apparent to those of ordinary skill in the art.

As detailed herein, the foregoing antibodies and other binding molecules may be used for example to identify tissues expressing protein or to purify protein. Antibodies also may be

coupled to specific diagnostic labeling agents for imaging of cells and tissues that express cancer associated antigens or to therapeutically useful agents according to standard coupling procedures. Diagnostic agents include, but are not limited to, barium sulfate, iocetamic acid, iopanoic acid, ipodate calcium, diatrizoate sodium, diatrizoate meglumine, metrizamide, tyropanoate sodium and radiodiagnostics including positron emitters such as fluorine-18 and carbon-11, gamma emitters such as iodine-123, technitium-99m, iodine-131 and indium-111, nuclides for nuclear magnetic resonance such as fluorine and gadolinium. Other diagnostic agents useful in the invention will be apparent to one of ordinary skill in the art. As used herein, "therapeutically useful agents" include any therapeutic molecule which desirably is targeted selectively to a cell expressing one of the cancer antigens disclosed herein, including antineoplastic agents, radioiodinated compounds, toxins, other cytostatic or cytolytic drugs, and so forth. Antineoplastic therapeutics are well known and include: aminoglutethimide, azathioprine, bleomycin sulfate, busulfan, carmustine, chlorambucil, cisplatin, cyclophosphamide, cyclosporine, cytarabidine, dacarbazine, dactinomycin, daunorubicin, doxorubicin, taxol, etoposide, fluorouracil, interferon- α , lomustine, mercaptopurine, methotrexate, mitotane, procarbazine HCl, thioguanine, vinblastine sulfate and vincristine sulfate. Additional antineoplastic agents include those disclosed in Chapter 52, Antineoplastic Agents (Paul Calabresi and Bruce A. Chabner), and the introduction thereto, 1202-1263, of Goodman and Gilman's "The Pharmacological Basis of Therapeutics", Eighth Edition, 1990, McGraw-Hill, Inc. (Health Professions Division). Toxins can be proteins such as, for example, pokeweed anti-viral protein, cholera toxin, pertussis toxin, ricin, gelonin, abrin, diphtheria exotoxin, or *Pseudomonas* exotoxin. Toxin moieties can also be high energy-emitting radionuclides such as cobalt-60.

In the foregoing methods, antibodies prepared according to the invention also preferably are specific for the cancer associated antigen/MHC complexes described herein.

When "disorder" is used herein, it refers to any pathological condition where the cancer associated antigens are expressed. An example of such a disorder is cancer, with fibrosarcoma as a particular example. For human cancers, additional particular examples include bladder cancer, breast cancer, lung cancer, colon cancer, and hepatoma.

Samples of tissue and/or cells for use in the various methods described herein can be obtained through standard methods such as tissue biopsy, including punch biopsy and cell scraping, and collection of blood or other bodily fluids by aspiration or other methods.

In certain embodiments of the invention, an immunoreactive cell sample is removed

from a subject. By "immunoreactive cell" is meant a cell which can mature into an immune cell (such as a B cell, a helper T cell, or a cytolytic T cell) upon appropriate stimulation. Thus immunoreactive cells include CD34⁺ hematopoietic stem cells, immature T cells and immature B cells. When it is desired to produce cytolytic T cells which recognize a cancer associated antigen, the immunoreactive cell is contacted with a cell which expresses a cancer associated antigen under conditions favoring production, differentiation and/or selection of cytolytic T cells; the differentiation of the T cell precursor into a cytolytic T cell upon exposure to antigen is similar to clonal selection of the immune system.

Some therapeutic approaches based upon the disclosure are premised on a response by a subject's immune system, leading to lysis of antigen presenting cells, such as cancer cells which present one or more cancer associated antigens. One such approach is the administration of autologous CTLs specific to a cancer associated antigen/MHC complex to a subject with abnormal cells of the phenotype at issue. It is within the ability of one of ordinary skill in the art to develop such CTLs *in vitro*. An example of a method for T cell differentiation is presented in International Application number PCT/US96/05607. Generally, a sample of cells taken from a subject, such as blood cells, are contacted with a cell presenting the complex and capable of provoking CTLs to proliferate. The target cell can be a transfectant, such as a COS cell. These transfectants present the desired complex of their surface and, when combined with a CTL of interest, stimulate its proliferation. COS cells are widely available, as are other suitable host cells. Specific production of CTL clones is well known in the art. The clonally expanded autologous CTLs then are administered to the subject.

Another method for selecting antigen-specific CTL clones has recently been described (Altman et al., *Science* 274:94-96, 1996; Dunbar et al., *Curr. Biol.* 8:413-416, 1998), in which fluorogenic tetramers of MHC class I molecule/peptide complexes are used to detect specific CTL clones. Briefly, soluble MHC class I molecules are folded *in vitro* in the presence of β_2 -microglobulin and a peptide antigen which binds the class I molecule. After purification, the MHC/peptide complex is purified and labeled with biotin. Tetramers are formed by mixing the biotinylated peptide-MHC complex with labeled avidin (e.g. phycoerythrin) at a molar ratio or 4:1. Tetramers are then contacted with a source of CTLs such as peripheral blood or lymph node. The tetramers bind CTLs which recognize the peptide antigen/MHC class I complex. Cells bound by the tetramers can be sorted by fluorescence activated cell sorting to isolate the reactive CTLs. The isolated CTLs then can be expanded *in vitro* for use as

described herein.

To detail a therapeutic methodology, referred to as adoptive transfer (Greenberg, *J. Immunol.* 136(5): 1917, 1986; Riddel et al., *Science* 257: 238, 1992; Lynch et al, *Eur. J. Immunol.* 21: 1403-1410,1991; Kast et al., *Cell* 59: 603-614, 1989), cells presenting the
5 desired complex (e.g., dendritic cells) are combined with CTLs leading to proliferation of the CTLs specific thereto. The proliferated CTLs are then administered to a subject with a cellular abnormality which is characterized by certain of the abnormal cells presenting the particular complex. The CTLs then lyse the abnormal cells, thereby achieving the desired therapeutic goal.

10 The foregoing therapy assumes that at least some of the subject's abnormal cells present the relevant HLA/cancer associated antigen complex. This can be determined very easily, as the art is very familiar with methods for identifying cells which present a particular HLA molecule, as well as how to identify cells expressing DNA of the pertinent sequences, in this case a cancer associated antigen sequence. Once cells presenting the relevant complex
15 are identified via the foregoing screening methodology, they can be combined with a sample from a patient, where the sample contains CTLs. If the complex presenting cells are lysed by the mixed CTL sample, then it can be assumed that a cancer associated antigen is being presented, and the subject is an appropriate candidate for the therapeutic approaches set forth *supra*.

20 Adoptive transfer is not the only form of therapy that is available in accordance with the invention. CTLs can also be provoked *in vivo*, using a number of approaches. One approach is the use of non-proliferative cells expressing the complex. The cells used in this approach may be those that normally express the complex, such as irradiated tumor cells or cells transfected with one or both of the genes necessary for presentation of the complex (i.e.
25 the antigenic peptide and the presenting MHC molecule). Chen et al. (*Proc. Natl. Acad. Sci. USA* 88: 110-114,1991) exemplifies this approach, showing the use of transfected cells expressing HPV E7 peptides in a therapeutic regime. Various cell types may be used. Similarly, vectors carrying one or both of the genes of interest may be used. Viral or bacterial vectors are especially preferred. For example, nucleic acids which encode a cancer associated
30 antigen polypeptide or peptide may be operably linked to promoter and enhancer sequences which direct expression of the cancer associated antigen polypeptide or peptide in certain tissues or cell types. The nucleic acid may be incorporated into an expression vector. Expression vectors may be unmodified extrachromosomal nucleic acids, plasmids or viral

genomes constructed or modified to enable insertion of exogenous nucleic acids, such as those encoding cancer associated antigen, as described elsewhere herein. Nucleic acids encoding a cancer associated antigen also may be inserted into a retroviral genome, thereby facilitating integration of the nucleic acid into the genome of the target tissue or cell type. In these systems, the gene of interest is carried by a microorganism, e.g., a Vaccinia virus, pox virus, herpes simplex virus, retrovirus or adenovirus, and the materials de facto "infect" host cells. The cells which result present the complex of interest, and are recognized by autologous CTLs, which then proliferate.

A similar effect can be achieved by combining the cancer associated antigen or a stimulatory fragment thereof with an adjuvant to facilitate incorporation into antigen presenting cells *in vivo*. The cancer associated antigen polypeptide is processed to yield the peptide partner of the MHC molecule while a cancer associated antigen peptide may be presented without the need for further processing. Generally, subjects can receive an intradermal injection of an effective amount of the cancer associated antigen. Initial doses can be followed by booster doses, following immunization protocols standard in the art. Preferred cancer associated antigens include those found to react with allogeneic cancer antisera, shown in the examples below.

The invention involves the use of various materials disclosed herein to "immunize" subjects or as "vaccines". As used herein, "immunization" or "vaccination" means increasing or activating an immune response against an antigen. It does not require elimination or eradication of a condition but rather contemplates the clinically favorable enhancement of an immune response toward an antigen. Generally accepted animal models, including the MethA fibroarcoma model used herein, can be used for testing of immunization against cancer using a cancer associated antigen nucleic acid. For example, human cancer cells can be introduced into a mouse to create a tumor, and one or more cancer associated antigen nucleic acids can be delivered by the methods described herein. The effect on the cancer cells (e.g., reduction of tumor size) can be assessed as a measure of the effectiveness of the cancer associated antigen nucleic acid immunization. Of course, testing of the foregoing animal model using more conventional methods for immunization include the administration of one or more cancer associated antigen polypeptides or peptides derived therefrom, optionally combined with one or more adjuvants and/or cytokines to boost the immune response. Methods for immunization, including formulation of a vaccine composition and selection of doses, route of administration and the schedule of administration (e.g. primary and one or more booster

doses), are well known in the art. The tests also can be performed in humans, where the end point is to test for the presence of enhanced levels of circulating CTLs against cells bearing the antigen, to test for levels of circulating antibodies against the antigen, to test for the presence of cells expressing the antigen and so forth.

5 As part of the immunization compositions, one or more cancer associated antigens or stimulatory fragments thereof are administered with one or more adjuvants to induce an immune response or to increase an immune response. An adjuvant is a substance incorporated into or administered with antigen which potentiates the immune response. Adjuvants may enhance the immunological response by providing a reservoir of antigen (extracellularly or
10 within macrophages), activating macrophages and stimulating specific sets of lymphocytes. Adjuvants of many kinds are well known in the art. Specific examples of adjuvants include monophosphoryl lipid A (MPL, SmithKline Beecham), a congener obtained after purification and acid hydrolysis of *Salmonella minnesota* Re 595 lipopolysaccharide; saponins including QS21 (SmithKline Beecham), a pure QA-21 saponin purified from *Quillja saponaria* extract;
15 DQS21, described in PCT application WO96/33739 (SmithKline Beecham); QS-7, QS-17, QS-18, and QS-L1 (So et al., *Mol. Cells* 7:178-186, 1997); incomplete Freund's adjuvant; complete Freund's adjuvant; montanide; alum; CpG oligonucleotides (see e.g. Kreig et al., *Nature* 374:546-9, 1995); and various water-in-oil emulsions prepared from biodegradable oils such as squalene and/or tocopherol. Preferably, the peptides are administered mixed with
20 a combination of DQS21/MPL. The ratio of DQS21 to MPL typically will be about 1:10 to 10:1, preferably about 1:5 to 5:1 and more preferably about 1:1. Typically for human administration, DQS21 and MPL will be present in a vaccine formulation in the range of about 1 µg to about 100 µg. Other adjuvants are known in the art and can be used in the invention (see, e.g. Goding, *Monoclonal Antibodies: Principles and Practice*, 2nd Ed., 1986).
25 Methods for the preparation of mixtures or emulsions of peptide and adjuvant are well known to those of skill in the art of vaccination.

Other agents which stimulate the immune response of the subject can also be administered to the subject. For example, other cytokines are also useful in vaccination protocols as a result of their lymphocyte regulatory properties. Many other cytokines useful
30 for such purposes will be known to one of ordinary skill in the art, including interleukin-12 (IL-12) which has been shown to enhance the protective effects of vaccines (see, e.g., *Science* 268: 1432-1434, 1995), GM-CSF and IL-18. Thus cytokines can be administered in conjunction with antigens and adjuvants to increase the immune response to the antigens.

There are a number of immune response potentiating compounds that can be used in vaccination protocols. These include costimulatory molecules provided in either protein or nucleic acid form. Such costimulatory molecules include the B7-1 and B7-2 (CD80 and CD86 respectively) molecules which are expressed on dendritic cells (DC) and interact with the CD28 molecule expressed on the T cell. This interaction provides costimulation (signal 2) to an antigen/MHC/TCR stimulated (signal 1) T cell, increasing T cell proliferation and effector function. B7 also interacts with CTLA4 (CD152) on T cells and studies involving CTLA4 and B7 ligands indicate that the B7-CTLA4 interaction can enhance antitumor immunity and CTL proliferation (Zheng P., et al. *Proc. Natl. Acad. Sci. USA* 95 (11):6284-6289 (1998)).

B7 typically is not expressed on tumor cells so they are not efficient antigen presenting cells (APCs) for T cells. Induction of B7 expression would enable the tumor cells to stimulate more efficiently CTL proliferation and effector function. A combination of B7/IL-6/IL-12 costimulation has been shown to induce IFN-gamma and a Th1 cytokine profile in the T cell population leading to further enhanced T cell activity (Gajewski et al., *J. Immunol.*, 154:5637-5648 (1995)). Tumor cell transfection with B7 has been discussed in relation to *in vitro* CTL expansion for adoptive transfer immunotherapy by Wang et al., (*J. Immunol.*, 19:1-8 (1986)). Other delivery mechanisms for the B7 molecule would include nucleic acid (naked DNA) immunization (Kim J., et al. *Nat. Biotechnol.*, 15:7:641-646 (1997)) and recombinant viruses such as adeno and pox (Wendtner et al., *Gene Ther.*, 4:7:726-735 (1997)). These systems are all amenable to the construction and use of expression cassettes for the coexpression of B7 with other molecules of choice such as the antigens or fragment(s) of antigens discussed herein (including polytopes) or cytokines. These delivery systems can be used for induction of the appropriate molecules *in vitro* and for *in vivo* vaccination situations. The use of anti-CD28 antibodies to directly stimulate T cells *in vitro* and *in vivo* could also be considered. Similarly, the inducible co-stimulatory molecule ICOS which induces T cell responses to foreign antigen could be modulated, for example, by use of anti-ICOS antibodies (Hutloff et al., *Nature* 397:263-266, 1999).

Lymphocyte function associated antigen-3 (LFA-3) is expressed on APCs and some tumor cells and interacts with CD2 expressed on T cells. This interaction induces T cell IL-2 and IFN-gamma production and can thus complement but not substitute, the B7/CD28 costimulatory interaction (Parra et al., *J. Immunol.*, 158:637-642 (1997), Fenton et al., *J. Immunother.*, 21:2:95-108 (1998)).

Lymphocyte function associated antigen-1 (LFA-1) is expressed on leukocytes and interacts with ICAM-1 expressed on APCs and some tumor cells. This interaction induces T cell IL-2 and IFN-gamma production and can thus complement but not substitute, the B7/CD28 costimulatory interaction (Fenton et al., *J. Immunother.*, 21:2:95-108 (1998)).

5 LFA-1 is thus a further example of a costimulatory molecule that could be provided in a vaccination protocol in the various ways discussed above for B7.

Complete CTL activation and effector function requires Th cell help through the interaction between the Th cell CD40L (CD40 ligand) molecule and the CD40 molecule expressed by DCs (Ridge et al., *Nature*, 393:474 (1998), Bennett et al., *Nature*, 393:478
10 (1998), Schoenberger et al., *Nature*, 393:480 (1998)). This mechanism of this costimulatory signal is likely to involve upregulation of B7 and associated IL-6/IL-12 production by the DC (APC). The CD40-CD40L interaction thus complements the signal 1 (antigen/MHC-TCR) and signal 2 (B7-CD28) interactions.

The use of anti-CD40 antibodies to stimulate DC cells directly, would be expected to
15 enhance a response to tumor antigens which are normally encountered outside of a inflammatory context or are presented by non-professional APCs (tumor cells). In these situations Th help and B7 costimulation signals are not provided. This mechanism might be used in the context of antigen pulsed DC based therapies or in situations where Th epitopes have not been defined within known TRA precursors.

20 A cancer associated antigen polypeptide, or a fragment thereof, also can be used to isolate their native binding partners. Isolation of such binding partners may be performed according to well-known methods. For example, isolated cancer associated antigen polypeptides can be attached to a substrate (e.g., chromatographic media, such as polystyrene beads, or a filter), and then a solution suspected of containing the binding partner may be
25 applied to the substrate. If a binding partner which can interact with cancer associated antigen polypeptides is present in the solution, then it will bind to the substrate-bound cancer associated antigen polypeptide. The binding partner then may be isolated.

It will also be recognized that the invention embraces the use of the cancer associated antigen cDNA sequences in expression vectors, as well as to transfect host cells and cell lines,
30 be these prokaryotic (e.g., *E. coli*), or eukaryotic (e.g., dendritic cells, B cells, CHO cells, COS cells, yeast expression systems and recombinant baculovirus expression in insect cells). Especially useful are mammalian cells such as human, mouse, hamster, pig, goat, primate, etc. They may be of a wide variety of tissue types, and include primary cells and cell lines.

Specific examples include dendritic cells, peripheral blood leukocytes, bone marrow stem cells and embryonic stem cells. The expression vectors require that the pertinent sequence, i.e., those nucleic acids described *supra*, be operably linked to a promoter.

The invention also contemplates delivery of nucleic acids, polypeptides or peptides for
5 vaccination. Delivery of polypeptides and peptides can be accomplished according to standard vaccination protocols which are well known in the art. In another embodiment, the delivery of nucleic acid is accomplished by *ex vivo* methods, i.e. by removing a cell from a subject, genetically engineering the cell to include a cancer associated antigen, and reintroducing the engineered cell into the subject. One example of such a procedure is
10 outlined in U.S. Patent 5,399,346 and in exhibits submitted in the file history of that patent, all of which are publicly available documents. In general, it involves introduction *in vitro* of a functional copy of a gene into a cell(s) of a subject, and returning the genetically engineered cell(s) to the subject. The functional copy of the gene is under operable control of regulatory elements which permit expression of the gene in the genetically engineered cell(s). Numerous
15 transfection and transduction techniques as well as appropriate expression vectors are well known to those of ordinary skill in the art, some of which are described in PCT application WO95/00654. *In vivo* nucleic acid delivery using vectors such as viruses and targeted liposomes also is contemplated according to the invention.

In preferred embodiments, a virus vector for delivering a nucleic acid encoding a
20 cancer associated antigen is selected from the group consisting of adenoviruses, adeno-associated viruses, poxviruses including vaccinia viruses and attenuated poxviruses, Semliki Forest virus, Venezuelan equine encephalitis virus, retroviruses, Sindbis virus, and Ty virus-like particle. Examples of viruses and virus-like particles which have been used to deliver exogenous nucleic acids include: replication-defective adenoviruses (e.g., Xiang et al.,
25 *Virology* 219:220-227, 1996; Eloit et al., *J. Virol.* 71:5375-5381, 1997; Chengalvala et al., *Vaccine* 15:335-339, 1997), a modified retrovirus (Townsend et al., *J. Virol.* 71:3365-3374, 1997), a nonreplicating retrovirus (Irwin et al., *J. Virol.* 68:5036-5044, 1994), a replication defective Semliki Forest virus (Zhao et al., *Proc. Natl. Acad. Sci. USA* 92:3009-3013, 1995), canarypox virus and highly attenuated vaccinia virus derivative (Paoletti, *Proc. Natl. Acad.*
30 *Sci. USA* 93:11349-11353, 1996), non-replicative vaccinia virus (Moss, *Proc. Natl. Acad. Sci. USA* 93:11341-11348, 1996), replicative vaccinia virus (Moss, *Dev. Biol. Stand.* 82:55-63, 1994), Venezuelan equine encephalitis virus (Davis et al., *J. Virol.* 70:3781-3787, 1996), Sindbis virus (Pugachev et al., *Virology* 212:587-594, 1995), and Ty virus-like particle

(Allsopp et al., *Eur. J. Immunol* 26:1951-1959, 1996). In preferred embodiments, the virus vector is an adenovirus.

Another preferred virus for certain applications is the adeno-associated virus, a double-stranded DNA virus. The adeno-associated virus is capable of infecting a wide range of cell types and species and can be engineered to be replication-deficient. It further has advantages, such as heat and lipid solvent stability, high transduction frequencies in cells of diverse lineages, including hematopoietic cells, and lack of superinfection inhibition thus allowing multiple series of transductions. The adeno-associated virus can integrate into human cellular DNA in a site-specific manner, thereby minimizing the possibility of insertional mutagenesis and variability of inserted gene expression. In addition, wild-type adeno-associated virus infections have been followed in tissue culture for greater than 100 passages in the absence of selective pressure, implying that the adeno-associated virus genomic integration is a relatively stable event. The adeno-associated virus can also function in an extrachromosomal fashion.

In general, other preferred viral vectors are based on non-cytopathic eukaryotic viruses in which non-essential genes have been replaced with the gene of interest. Non-cytopathic viruses include retroviruses, the life cycle of which involves reverse transcription of genomic viral RNA into DNA with subsequent proviral integration into host cellular DNA. Adenoviruses and retroviruses have been approved for human gene therapy trials. In general, the retroviruses are replication-deficient (i.e., capable of directing synthesis of the desired proteins, but incapable of manufacturing an infectious particle). Such genetically altered retroviral expression vectors have general utility for the high-efficiency transduction of genes *in vivo*. Standard protocols for producing replication-deficient retroviruses (including the steps of incorporation of exogenous genetic material into a plasmid, transfection of a packaging cell lined with plasmid, production of recombinant retroviruses by the packaging cell line, collection of viral particles from tissue culture media, and infection of the target cells with viral particles) are provided in Kriegler, M., "Gene Transfer and Expression, A Laboratory Manual," W.H. Freeman Co., New York (1990) and Murry, E.J. Ed. "Methods in Molecular Biology," vol. 7, Humana Press, Inc., Clifton, New Jersey (1991).

Preferably the foregoing nucleic acid delivery vectors: (1) contain exogenous genetic material that can be transcribed and translated in a mammalian cell and that can induce an immune response in a host, and (2) contain on a surface a ligand that selectively binds to a receptor on the surface of a target cell, such as a mammalian cell, and thereby gains entry to

the target cell.

Various techniques may be employed for introducing nucleic acids of the invention into cells, depending on whether the nucleic acids are introduced *in vitro* or *in vivo* in a host. Such techniques include transfection of nucleic acid-CaPO₄ precipitates, transfection of
5 nucleic acids associated with DEAE, transfection or infection with the foregoing viruses including the nucleic acid of interest, liposome mediated transfection, and the like. For certain uses, it is preferred to target the nucleic acid to particular cells. In such instances, a vehicle used for delivering a nucleic acid of the invention into a cell (e.g., a retrovirus, or other virus; a liposome) can have a targeting molecule attached thereto. For example, a
10 molecule such as an antibody specific for a surface membrane protein on the target cell or a ligand for a receptor on the target cell can be bound to or incorporated within the nucleic acid delivery vehicle. Preferred antibodies include antibodies which selectively bind a cancer associated antigen, alone or as a complex with a MHC molecule. Especially preferred are monoclonal antibodies. Where liposomes are employed to deliver the nucleic acids of the
15 invention, proteins which bind to a surface membrane protein associated with endocytosis may be incorporated into the liposome formulation for targeting and/or to facilitate uptake. Such proteins include capsid proteins or fragments thereof tropic for a particular cell type, antibodies for proteins which undergo internalization in cycling, proteins that target intracellular localization and enhance intracellular half life, and the like. Polymeric delivery
20 systems also have been used successfully to deliver nucleic acids into cells, as is known by those skilled in the art. Such systems even permit oral delivery of nucleic acids.

When administered, the therapeutic compositions of the present invention can be administered in pharmaceutically acceptable preparations. Such preparations may routinely contain pharmaceutically acceptable concentrations of salt, buffering agents, preservatives,
25 compatible carriers, supplementary immune potentiating agents such as adjuvants and cytokines and optionally other therapeutic agents.

The therapeutics of the invention can be administered by any conventional route, including injection or by gradual infusion over time. The administration may, for example, be oral, intravenous, intraperitoneal, intramuscular, intracavity, subcutaneous, or transdermal.
30 When antibodies are used therapeutically, a preferred route of administration is by pulmonary aerosol. Techniques for preparing aerosol delivery systems containing antibodies are well known to those of skill in the art. Generally, such systems should utilize components which will not significantly impair the biological properties of the antibodies, such as the paratope

binding capacity (see, for example, Sciarra and Cutie, "Aerosols," in Remington's Pharmaceutical Sciences, 18th edition, 1990, pp 1694-1712; incorporated by reference).

Those of skill in the art can readily determine the various parameters and conditions for producing antibody aerosols without resort to undue experimentation. When using antisense preparations of the invention, slow intravenous administration is preferred.

The compositions of the invention are administered in effective amounts. An "effective amount" is that amount of a cancer associated antigen composition that alone, or together with further doses, produces the desired response, e.g. increases an immune response to the cancer associated antigen. In the case of treating a particular disease or condition characterized by expression of one or more cancer associated antigens, such as cancer, the desired response is inhibiting the progression of the disease. This may involve only slowing the progression of the disease temporarily, although more preferably, it involves halting the progression of the disease permanently. This can be monitored by routine methods or can be monitored according to diagnostic methods of the invention discussed herein. The desired response to treatment of the disease or condition also can be delaying the onset or even preventing the onset of the disease or condition.

Such amounts will depend, of course, on the particular condition being treated, the severity of the condition, the individual patient parameters including age, physical condition, size and weight, the duration of the treatment, the nature of concurrent therapy (if any), the specific route of administration and like factors within the knowledge and expertise of the health practitioner. These factors are well known to those of ordinary skill in the art and can be addressed with no more than routine experimentation. It is generally preferred that a maximum dose of the individual components or combinations thereof be used, that is, the highest safe dose according to sound medical judgment. It will be understood by those of ordinary skill in the art, however, that a patient may insist upon a lower dose or tolerable dose for medical reasons, psychological reasons or for virtually any other reasons.

The pharmaceutical compositions used in the foregoing methods preferably are sterile and contain an effective amount of cancer associated antigen or nucleic acid encoding cancer associated antigen for producing the desired response in a unit of weight or volume suitable for administration to a patient. The response can, for example, be measured by determining the immune response following administration of the cancer associated antigen composition via a reporter system by measuring downstream effects such as gene expression, or by measuring the physiological effects of the cancer associated antigen composition, such as

regression of a tumor or decrease of disease symptoms. Other assays will be known to one of ordinary skill in the art and can be employed for measuring the level of the response.

The doses of cancer associated antigen compositions (e.g., polypeptide, peptide, antibody, cell or nucleic acid) administered to a subject can be chosen in accordance with different parameters, in particular in accordance with the mode of administration used and the state of the subject. Other factors include the desired period of treatment. In the event that a response in a subject is insufficient at the initial doses applied, higher doses (or effectively higher doses by a different, more localized delivery route) may be employed to the extent that patient tolerance permits.

In general, for treatments for eliciting or increasing an immune response, doses of cancer associated antigen are formulated and administered in doses between 1 ng and 1 mg, and preferably between 10 ng and 100 µg, according to any standard procedure in the art. Where nucleic acids encoding cancer associated antigen of variants thereof are employed, doses of between 1 ng and 0.1 mg generally will be formulated and administered according to standard procedures. Other protocols for the administration of cancer associated antigen compositions will be known to one of ordinary skill in the art, in which the dose amount, schedule of injections, sites of injections, mode of administration (e.g., intra-tumoral) and the like vary from the foregoing. Administration of cancer associated antigen compositions to mammals other than humans, e.g. for testing purposes or veterinary therapeutic purposes, is carried out under substantially the same conditions as described above.

Where cancer associated antigen peptides are used for vaccination, modes of administration which effectively deliver the cancer associated antigen and adjuvant, such that an immune response to the antigen is increased, can be used. For administration of a cancer associated antigen peptide in adjuvant, preferred methods include intradermal, intravenous, intramuscular and subcutaneous administration. Although these are preferred embodiments, the invention is not limited by the particular modes of administration disclosed herein. Standard references in the art (e.g., *Remington's Pharmaceutical Sciences*, 18th edition, 1990) provide modes of administration and formulations for delivery of immunogens with adjuvant or in a non-adjuvant carrier.

When administered, the pharmaceutical preparations of the invention are applied in pharmaceutically-acceptable amounts and in pharmaceutically-acceptable compositions. The term "pharmaceutically acceptable" means a non-toxic material that does not interfere with the effectiveness of the biological activity of the active ingredients. Such preparations may

routinely contain salts, buffering agents, preservatives, compatible carriers, and optionally other therapeutic agents. When used in medicine, the salts should be pharmaceutically acceptable, but non-pharmaceutically acceptable salts may conveniently be used to prepare pharmaceutically-acceptable salts thereof and are not excluded from the scope of the

5 invention. Such pharmacologically and pharmaceutically-acceptable salts include, but are not limited to, those prepared from the following acids: hydrochloric, hydrobromic, sulfuric, nitric, phosphoric, maleic, acetic, salicylic, citric, formic, malonic, succinic, and the like. Also, pharmaceutically-acceptable salts can be prepared as alkaline metal or alkaline earth salts, such as sodium, potassium or calcium salts.

10 A cancer associated antigen composition may be combined, if desired, with a pharmaceutically-acceptable carrier. The term "pharmaceutically-acceptable carrier" as used herein means one or more compatible solid or liquid fillers, diluents or encapsulating substances which are suitable for administration into a human. The term "carrier" denotes an organic or inorganic ingredient, natural or synthetic, with which the active ingredient is
15 combined to facilitate the application. The components of the pharmaceutical compositions also are capable of being co-mingled with the molecules of the present invention, and with each other, in a manner such that there is no interaction which would substantially impair the desired pharmaceutical efficacy.

The pharmaceutical compositions may contain suitable buffering agents, including:
20 acetic acid in a salt; citric acid in a salt; boric acid in a salt; and phosphoric acid in a salt.

The pharmaceutical compositions also may contain, optionally, suitable preservatives, such as: benzalkonium chloride; chlorobutanol; parabens and thimerosal.

The pharmaceutical compositions may conveniently be presented in unit dosage form and may be prepared by any of the methods well-known in the art of pharmacy. All methods
25 include the step of bringing the active agent into association with a carrier which constitutes one or more accessory ingredients. In general, the compositions are prepared by uniformly and intimately bringing the active compound into association with a liquid carrier, a finely divided solid carrier, or both, and then, if necessary, shaping the product.

Compositions suitable for oral administration may be presented as discrete units, such
30 as capsules, tablets, lozenges, each containing a predetermined amount of the active compound. Other compositions include suspensions in aqueous liquids or non-aqueous liquids such as a syrup, elixir or an emulsion.

Compositions suitable for parenteral administration conveniently comprise a sterile

aqueous or non-aqueous preparation of cancer associated antigen polypeptides or nucleic acids, which is preferably isotonic with the blood of the recipient. This preparation may be formulated according to known methods using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation also may be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example, as a solution in 1,3-butane diol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono-or di-glycerides. In addition, fatty acids such as oleic acid may be used in the preparation of injectables. Carrier formulation suitable for oral, subcutaneous, intravenous, intramuscular, etc. administrations can be found in *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, PA.

Examples

Example 1: Identification of Mouse Tumor Antigens by SEREX

To investigate tumor specificity in host immune reactions to SEREX antigens, the well-controlled mouse tumor model BALB/c methylcholanthrene-induced fibrosarcoma Meth A was used. Meth A is a prototype tumor for analysis of tumor rejection antigens (Gross, L., *Cancer Res.* 3:326-333, 1943; Foley, *Cancer Res.* 13:835-842, 1953; Prehn et al., *J. Natl. Cancer Inst.* 18:769-778, 1957; and Old et al., *Ann. N.Y. Acad. Sci.* 101:80-106, 1962).

Materials and Methods

Mice

BALB/c mice were purchased from Japan SLC (Shizuoka, Japan).

Tumors and sera

Meth A, CMS5a, CMS5j, CMS8, CMS13 and CMS17 are methylcholanthrene-induced fibrosarcomas in BALB/c mice (DeLeo et al., *J. Exp. Med.* 146:720-734, 1977; Palladino et al., *Cancer Res.* 47:5074-5079, 1987). RL σ 1 is a radiation-induced leukemia in a BALB/c mouse (Nakayama et al., *Proc. Natl. Acad. Sci USA* 76:3486-3490, 1979). RVA, RVC and RVD are leukemias induced by an injection of radiation leukemia virus into neonatal BALB/c mice (Stockert et al., *J. Exp. Med.* 149:200-215, 1984). RV2 is leukemia induced by an injection of radiation leukemia virus into a neonatal C57BL/6 mouse

(Nakayama et al., *Cancer Res.* 44:5138-5144, 1984). A20.2J is a BALB/c B cell lymphoma (Kim et al., *J. Immunol.* 122:549-554, 1979). MOPC-70A is a mineral oil-induced myeloma in a BALB/c mouse (Potter, M. In: *Methods in Cancer Research*, 1967, pp. 106-157). P815 is a methylcholanthrene-induced mastocytoma in a DBA/2 mouse (Dunn et al., *J. Natl. Cancer*
5 *Inst.* 18:587-601, 1957). EL4 is an dimethylbenzanthracene-induced leukemia in a C57BL mouse (Gorer et al., *Br. J. Cancer* 4:3720379, 1950).

Meth A (K γ) cells used for construction of cDNA expression library were Meth A cells which were retrovirally introduced with murine IFN- γ cDNA (Watanabe et al., *Eur. J. Immunol.* 18:1627-1630, 1988). Meth A (K γ) cells were highly antigenic and regressed in
10 CB6F₁ mice. Concomitant inoculation of the cells caused rejection of the parental Meth A cells (Harutsumi et al., *Int. J. Oncol.* 7:233-238, 1995).

Sera were obtained from BALB/c mice at various days after 5 x 10⁵ Meth A inoculation (i.d. in the back). Sera were obtained from 3-5 mice each at 16, 23, 31 and 45 days after MethA inoculation.

15 *Construction of cDNA Expression Library.*

Poly (A)⁺ RNA was purified from Meth A (K γ) cells and a cDNA expression library was constructed in a λ ZAP Express vector using a cDNA library kit (Stratagene, La Jolla, CA).

20 *Immunoscreening of cDNA Library.*

cDNA libraries was screened with serum from BALB/c mice obtained on day 45 after inoculation of Meth A. The nitrocellulose membranes containing the phages were incubated overnight at room temperature with the serum diluted to 1:200, which has been preabsorbed
25 with lysate from *E. coli* coupled to Sepharose 4B (5 Prime \rightarrow 3 Prime, Boulder, CO). Serum antibodies binding to recombinant proteins expressed in lytic phages were detected by incubation with peroxidase-conjugated goat anti-mouse IgG (Jackson Immuno Research, West Grove, PA) and visualized by staining with 3-3'-diaminobenzidine (Sigma, St. Louis, MO).

30 *Sequence Analysis of Reactive Clones.*

The reactive clones were subcloned to monoclonality and excised *in vivo* to pBK-CMV plasmid forms (Stratagene, La Jolla, CA). The nucleotide sequence of cDNA inserts

was determined by using ABI PRISM automated sequencers (PE Applied Biosystems, Foster City, CA). Sequence alignments were performed with BLAST software on GenBank.

Analysis of antigenicity of SEREX antigens.

To analyze antigenicity of SEREX antigens, sera from BALB/c mice bearing CMS5a, CMS5j, CMS8, CMS13 and CMS17 were used for the detection of antibodies with the immunoscreening assay described above. Serum from BALB/c mice obtained 42 weeks s.c injection with 3-methylcholanthrene dissolved in peanut oil at a concentration of 0.125 µg/ml was also used.

Reverse transcription PCR (RT-PCR)

mRNA was purified from normal BALB/c mouse tissues and tumors using the QuickPrep Micro mRNA Purification Kit (Amersham Pharmacia, Piscataway, NJ). The mRNA was reverse transcribed into single strand cDNA using Moloney murine leukemia virus reverse transcriptase and oligo(DT)₁₅ as primer (Amersham Pharmacia, Piscataway, NJ). cDNAs were tested for integrity by amplification of β-actin transcripts in a 30 cycle reaction. Gene specific oligonucleotide primers were designed to amplify cDNA segments for 300-600 bp in length. RT-PCR was performed by using 30 amplification cycles at an annealing temperature of 58°C using the following primers:

MC4A 5'-GTGGACAAGAGGAAGCACAA-3' (SEQ ID NO:21) and
MC4B 5'-TGAAAAGTAAGGGCTGTCAT-3' (SEQ ID NO:22).

Isolation and characterization of antigens in Meth A by SEREX

cDNA expression libraries were prepared from Meth A (Kγ) cells, which were Meth A cells retrovirally introduced with IFN-γ in order to augment antigenicity. Immunoscreening of 80,000 clones with serum from BALB/c mice obtained at day 45 after inoculation of parental Meth A yielded a total of 35 positive clones. The nucleotide sequences of the cDNA inserts were determined. As shown in Table 1, ten different genes were isolated (OY-MC-1 through OY-MC-10), five of which were identical to or had homology to genes in the DNA database. Of these, OY-MC-1 was represented by 13 overlapping clones, OY-MC-2 by 10 overlapping clones, OY-MC-3 by 3 overlapping clones, OY-MC-4 by 2 overlapping clones and OY-MC-6 by a single clone. OY-MC-1 was homologous to rat ribosomal protein L11. OY-MC-3 was homologous to bovine NAD⁺ dependent isocitrate dehydrogenase subunits 3

and 4. OY-MC-2 was homologous to mouse glyceraldehyde-3-phosphate dehydrogenase (GAPDH). OY-MC-4 was homologous to mouse placenta and embryonic expression gene (*pem*). OY-MC-6 was identical to a C4 and cytochrome P450 hydroxylase A gene. The remaining five genes (OY-MC-5, 7, 8, 9, 10) were unknown genes which had no homology to any published genes.

Table 1. Genes Isolated from Meth A by SEREX

Gene	Size, bp	SEQ ID NO	Homology/Identity	Frequency
OY-MC-1	585	1	<i>R. rattus</i> ribosomal protein L11	13/35
OY-MC-2	1228	3	mouse glyceraldehyde-3-phosphate dehydrogenase	10/35
OY-MC-3	1758	5	<i>B. taurus</i> NAD ⁺ dependent isocitrate dehydrogenase subunits 3 and 4	3/35
OY-MC-4	845	7	mouse placenta and embryonic expression gene (<i>pem</i>)	2/35
OY-MC-5	~1,000	9	No strong homology	2/35
OY-MC-6	~2,100	11	C4 and cytochrome P450 hydroxylase A	1/35
OY-MC-7	~1,400	13	No strong homology	1/35
OY-MC-8	612	15	No strong homology	1/35
OY-MC-9	1,036	17	No strong homology	1/35
OY-MC-10	~500	19	No strong homology	1/35

Expression of SEREX-defined antigens in normal tissues and tumors

mRNA expression for the genes were examined by RT-PCR, using a panel of normal mouse tissues and tumors. OY-MC-1, OY-MC-2 and OY-MC-9 showed ubiquitous mRNA expression in normal tissues, other genes showed a variety of expression patterns. Of those, OY-MC-4 showed a restricted expression in normal tissues as it was expressed only in testis and placenta, and to a lesser extent in ovary (Table 2).

Table 2. mRNA Expression of OY-MC-4 in Normal Mouse Tissues

Tissue	MRNA	Tissue	MRNA
Brain	-	Kidney	-
Fetal brain	-	Pancreas	-
Salivary gland	-	Stomach	-
Esophagus	-	Duodenum	-
Thymus	-	Small intestine	-
Heart	-	Colon	-
Lung	-	Bladder	-
Liver	-	Uterus	-
Fetal liver	-	Ovary	+#
Spleen	-	Testis	+
Lymph node	-	Placenta	+
Adrenal gland	-	Bone marrow	-

[#]Weak expression

Nine out of ten genes except OY-MC-4 showed ubiquitous expression in 15 mouse tumors tested. Expression of OY-MC-4 was rather restricted and found in 10 of 15 mouse tumors tested (Table 3). Of 6 methylcholanthrene-induced fibrosarcomas examined, Meth A, CMS5j, CMS8 and CMS13 showed OY-MC-4 expression, CMS17 showed a faint expression, and CMS5a was negative. Expression of OY-MC-4 was much less frequent in 6 leukemias examined. Strong expression of OY-MC-4 was only found in EL4. RVD and RV2 showed a faint expression, and RL ♂ 1, RVA, RVC were negative (Table 3).

Table 3. mRNA Expression of OY-MC-4 in Mouse Tumors

Tumor	mRNA	Tumor	mRNA
Meth A	+	RVC	-
CMS5a	-	RVD	± [#]
CMS5j	+	RV2	± [#]
CMS8	+	EL4	+
CMS13	+	A20.2j	+
CMS17	± [#]	MOPC-70A	-
RL ♂ 1	-	P815	+
RVA	-		

[#] Faint expression.

cDNA and protein sequence of OY-MC-4

The nucleotide sequence of OY-MC-4 revealed that the cDNA was 845 bp in length (SEQ ID NO:7) and nearly identical to a mouse homeobox gene, placenta and embryonic expression gene (*pem*). The *pem* gene was identified from an AKR T-cell lymphoma and shown to be coded for by the X chromosome (Wilkinson et al., *Dev. Biol.* 141:451-455, 1990; Lin et al., *Dev. Biol.* 166:170-179, 1994). One nucleotide difference (C to A) at position 619 was observed in OY-MC-4 compared with the published *pem* sequence. This nucleotide difference results in a Thr→Asp change at amino acid residue 174 (SEQ ID NO:8) as compared with the published *pem* sequence. The sequences of OY-MC-4 PCR products from Meth A, CMS5j, CMS8, CMS13 and testis that were subcloned into pCR2.1 showed A (adenine) at position 619, indicating that A at position 619 of SEQ ID NO:7 is a normal *pem* nucleotide sequence in BALB/c mouse.

Reactivity of sera of various stages of Meth A bearing mice

To determine the occurrence of antibodies against the defined antigens, sera obtained at various stages from parental Meth A bearing BALB/c mice were tested. As shown in Table

4, antibodies against 7 out of 10 antigens were detected in sera from mice at day 31 after inoculation of Meth A, and increased in titer at day 45. OY-MC-4, OY-MC-6 and OY-MC-8 were only detected by sera obtained on day 45.

5 **Table 4. Reaction of Serum Obtained From Various Stages of Meth A Bearing Mice**

Gene	Days After Tumor Inoculation			
	16d	23d	31d	45d
OY-MC-1	-	-	+	+
OY-MC-2	-	±	+	+
OY-MC-3	-	+	+	+
OY-MC-4	-	-	±	+
OY-MC-5	-	-	+	+
OY-MC-6	-	-	±	+
OY-MC-7	-	-	+	+
OY-MC-8	-	-	-	+
OY-MC-9	-	-	+	+
OY-MC-10	-	-	+	+

Sera obtained from mice at day 45 after Meth A inoculation showed the strongest reactivities against all of the antigens compares with those from mice at day 31 and 23 after Meth A inoculation.

10

Antigenicity of SEREX defined antigens

To analyze antigenicity of SEREX-defined antigens, sera from BALB/c mice bearing a panel of methylcholanthrene sarcomas were studied. As shown in Table 5, all the antigens were only detected with sera from the parental Meth A bearing mice.

15

Table 5. Specificity of SEREX antigens

	Tumor					
	Meth A	CMS 5a#	CMS 5j#	CMS 8#	CMS 13#	Methylcholanthrene*
OY-MC-1	+	-	-	-	-	-
OY-MC-2	+	-	-	-	-	-
OY-MC-3	+	-	-	-	-	-
OY-MC-4	+	-	-	-	-	-
OY-MC-5	+	-	-	-	-	-
OY-MC-6	+	-	-	-	-	-
OY-MC-7	+	-	-	-	-	-
OY-MC-8	+	-	-	-	-	-
OY-MC-9	+	-	-	-	-	-
OY-MC-10	+	-	-	-	-	-

Methylcholanthrene-induced fibrosarcomas.

* BALB/c mice were injected s.c. with 3-methylcholanthrene dissolved in peanut oil at a concentration of 125 µg/ml in a volume of 0.2 ml. Serum was obtained 42 weeks after injection.

5

Ten different genes were identified by immunoscreening of a cDNA expression library from Meth A (K γ) cells with sera from parental Meth A bearing BALB/c mice. Of those 10 genes, 5 were identical to or had homology to genes in the DNA database, and 5 were unknown genes. RT-PCR analysis revealed a variety of mRNA expression patterns of the genes in normal mouse tissues. Of most interest was OY-MC-4, which showed a restricted expression in normal tissues. It was strongly expressed in testis and placenta, and to a lesser extent in ovary. No expression was found in the other 21 out of 24 normal tissues tested. Furthermore, OY-MC-4 was selectively expressed in a variety of mouse tumors. Of 6 methylcholanthrene-induced fibrosarcomas examined, 4 (Meth A, CMS5j, CMS8, CMS13) showed OY-MC-4 expression, one (CMS17) showed a faint expression, whereas CMS5a was negative. OY-MC-4 expression in leukemias was much less frequent than those in fibrosarcomas: only EL4 showed strong expression, other five were negative or showed a faint expression.

Nucleotide sequencing analysis revealed that OY-MC-4 was nearly identical to mouse placenta and embryonic expression gene (*pem*), which was previously shown to be expressed during embryogenesis (Wilkinson et al., *Dev. Biol.* 141:451-455, 1990). *Pem* gene was distantly related to the *prd/pax* homeobox gene family (Lin et al., *Dev. Biol.* 166:170-179, 1994; Sasaki et al., *Mech. Dev.* 34:155-164, 1991) and was shown to be coded for by the X chromosome (Lin et al., *Dev. Biol.* 166:170-179, 1994; Maiti et al., *Genomics* 34:304-316, 1996). OY-MC-4 nucleotide sequence (SEQ ID NO:7) differed from the published *pem* cDNA sequence, which was isolated from a T-lymphoma of AKR mouse, in having an A instead of a C at position 619. The corresponding predicted amino acid sequence of OY-MC-4 (SEQ ID NO:8) included asparagine instead of threonine at position 174. Threonine at position 174 was thought to be a phosphorylation site for protein kinase C (Wilkinson et al., 1990. *Dev. Biol.* 141:451-455). OY-MC-4 cDNAs isolated from other tumors (CMS5j, CMS8, CMS13) and normal BALB/c testis had A at position 619, indicating that the sequence isolated from Meth A is not a point mutation but a normal nucleotide sequence of *pem* in BALB/c mouse. Taken together, the findings suggested that OY-MC-4 is a new cancer/testis (CT) antigen isolated by the SEREX method.

There are a growing number of genes belonging to the CT antigen family, including

MAGE, BAGE, GAGE, HOM-MEL-40/SSX2, NY-ESO-1, SCP1 and CT7. Of those, MAGE and NY-ESO-1 have been shown to elicit both humoral and cytotoxic T lymphocyte responses. NY-ESO-1 was isolated from an esophageal squamous cell carcinoma. NY-ESO-1 has been shown to be recognized by CTL, in the context of HLA-A2 and HLA-A31. SCP1, which was isolated from a cDNA library enriched for testis specific clones by using sera from patients with renal cancer, is presently the only CT antigen with known function. The SCP1 protein is involved in the meiotic chromosome synapses of sperm cells. OY-MC-4, described here, is a new member of SEREX-defined CT antigens with known function.

10 **Example 2: Preparation of recombinant cancer associated antigens**

To facilitate screening of mice or patients' sera for antibodies reactive with cancer associated antigens, for example by ELISA, recombinant proteins are prepared according to standard procedures. In one method, the clones encoding cancer associated antigens are subcloned into a baculovirus expression vector, and the recombinant expression vectors are introduced into appropriate insect cells. Baculovirus/insect cloning systems are preferred because post-translational modifications are carried out in the insect cells. Another preferred eukaryotic system is the *Drosophila* Expression System from Invitrogen. Clones which express high amounts of the recombinant protein are selected and used to produce the recombinant proteins. The recombinant proteins are tested for antibody recognition using serum from the mouse or patient which was used to isolated the particular clone, or in the case of cancer associated antigens recognized by allogeneic sera, by the sera from any of the mouse strains or patients used to isolate the clones or sera which recognize the clones' gene products.

Alternatively, the cancer associated antigen clones are inserted into a prokaryotic expression vector for production of recombinant proteins in bacteria. Other systems, including yeast expression systems and mammalian cell culture systems also can be used.

Example 3: Preparation of antibodies to cancer associated antigens

The recombinant cancer associated antigens produced as in Example 2 above are used to generate polyclonal antisera and monoclonal antibodies according to standard procedures. The antisera and antibodies so produced are tested for correct recognition of the cancer associated antigens by using the antisera/antibodies in assays of cell extracts of mice or patients known to express the particular cancer associated antigen (e.g. an ELISA assay). These antibodies can be used for experimental purposes (e.g. localization of the cancer

associated antigens, immunoprecipitations, Western blots, etc.) as well as diagnostic purposes (e.g., testing extracts of tissue biopsies, testing for the presence of cancer associated antigens).

Example 4: Expression of cancer associated antigens in cancers of similar and different origin.

The expression of one or more of the cancer associated antigens is tested in a range of tumor samples to determine which, if any, other malignancies should be diagnosed and/or treated by the methods described herein. Tumor cell lines and tumor samples are tested for cancer associated antigen expression, preferably by RT-PCR according to standard procedures. Northern blots also are used to test the expression of the cancer associated antigens. Antibody based assays, such as ELISA and western blot, also can be used to determine protein expression. A preferred method of testing expression of cancer associated antigens (in other cancers and in additional same type cancer patients or mice) is allogeneic serotyping using a modified SEREX protocol (as described above).

In all of the foregoing, extracts from the tumors of mice or patients who provided sera for the initial isolation of the cancer associated antigens are used as positive controls. The cells containing recombinant expression vectors described in the Examples above also can be used as positive controls.

The results generated from the foregoing experiments provide panels of multiple cancer associated nucleic acids and/or polypeptides for use in diagnostic (e.g. determining the existence of cancer, determining the prognosis of a patient undergoing therapy, etc.) and therapeutic methods (e.g., vaccine composition, etc.).

Example 5: HLA typing of patients positive for cancer associated antigen

To determine which HLA molecules present peptides derived from the cancer associated antigens, cells of the patients which express the cancer associated antigens are HLA typed (mice likewise can be typed for relevant MHC molecules). Peripheral blood lymphocytes are taken from the patient and typed for HLA class I or class II, as well as for the particular subtype of class I or class II. Tumor biopsy samples also can be used for typing. HLA typing can be carried out by any of the standard methods in the art of clinical immunology, such as by recognition by specific monoclonal antibodies, or by HLA allele-specific PCR (e.g. as described in WO97/31126).

Example 6: Characterization of cancer associated antigen peptides presented by MHC class I and class II molecules.

Antigens which provoke an antibody response in a subject may also provoke a cell-mediated immune response. Cells process proteins into peptides for presentation on MHC class I or class II molecules on the cell surface for immune surveillance. Peptides presented by certain MHC/HLA molecules generally conform to motifs. These motifs are known in some cases, and can be used to screen the cancer associated antigens for the presence of potential class I and/or class II peptides. Summaries of class I and class II motifs have been published (e.g., Rammensee et al., *Immunogenetics* 41:178-228, 1995). Based on the results of experiments such as those described above, the HLA types which present the individual cancer associated antigens are known. Motifs of peptides presented by these HLA molecules thus are preferentially searched.

One also can search for class I and class II motifs using computer algorithms. For example, computer programs for predicting potential CTL epitopes based on known class I motifs has been described (see, e.g., Parker et al, *J. Immunol.* 152:163, 1994; D'Amato et al., *Human Immunol.* 43:13-18, 1995; Drijfhout et al., *Human Immunol.* 43:1-12, 1995). HLA binding predictions can conveniently be made using an algorithm available via the Internet on the National Institutes of Health World Wide Web site at URL <http://bimas.dcrt.nih.gov>. See also the website of: SYFPEITHI: An Internet Database for MHC Ligands and Peptide Motifs (access via <http://www.uni-tuebingen.de/uni/kxi/> or <http://134.2.96.221/scripts/hlaserver.dll/EpPredict.htm>. Methods for determining HLA class II peptides and making substitutions thereto are also known (e.g. Strominger and Wucherpennig (PCT/US96/03182)).

Example 7: Identification of the portion of a cancer associated polypeptide encoding an antigen

To determine if the cancer associated antigens isolated as described above can provoke a cytolytic T lymphocyte response, the following method is performed. CTL clones are generated by stimulating the peripheral blood lymphocytes (PBLs) of a patient (or mouse) with autologous normal cells transfected with one of the clones encoding a cancer associated antigen polypeptide or with irradiated PBLs loaded with synthetic peptides corresponding to the putative protein and matching the consensus for the appropriate HLA class I molecule (as described above) to localize an antigenic peptide within the cancer associated antigen clone

(see, e.g., Knuth et al., *Proc. Natl. Acad. Sci. USA* 81:3511-3515, 1984; van der Bruggen et al., *Eur. J. Immunol.* 24:3038-3043, 1994). These CTL clones are screened for specificity against COS cells transfected with the cancer associated antigen clone and autologous HLA alleles as described by Brichard et al. (*Eur. J. Immunol.* 26:224-230, 1996). CTL recognition of a cancer associated antigen is determined by measuring release of TNF from the cytolytic T lymphocyte or by ^{51}Cr release assay (Herin et al., *Int. J. Cancer* 39:390-396, 1987). If a CTL clone specifically recognizes a transfected COS cell, then shorter fragments of the cancer associated antigen clone transfected in that COS cell are tested to identify the region of the gene that encodes the peptide. Fragments of the cancer associated antigen clone are prepared by exonuclease III digestion or other standard molecular biology methods. Synthetic peptides are prepared to confirm the exact sequence of the antigen.

Optionally, shorter fragments of cancer associated antigen cDNAs are generated by PCR. Shorter fragments are used to provoke TNF release or ^{51}Cr release as above.

Synthetic peptides corresponding to portions of the shortest fragment of the cancer associated antigen clone which provokes TNF release are prepared. Progressively shorter peptides are synthesized to determine the optimal cancer associated antigen tumor rejection antigen peptides for a given HLA molecule.

A similar method is performed to determine if the cancer associated antigen contains one or more HLA class II peptides recognized by T cells. One can search the sequence of the cancer associated antigen polypeptides for HLA class II motifs as described above. In contrast to class I peptides, class II peptides are presented by a limited number of cell types. Thus for these experiments, dendritic cells or B cell clones which express HLA class II molecules preferably are used.

Example 8: Identification of acrosomal protein, sp32, as a new human cancer/testis antigen.

RNA isolation and cDNA synthesis

Total RNA was isolated from human testis using the RNeasy kit (Qiagen GmbH, Germany). Poly (A)⁺ RNA was isolated from tumor tissue using the QuickPrep Micro mRNA Purification kit (Amersham Pharmacia, Piscataway, NJ). The mRNA was reverse transcribed into single strand cDNA using the Moloney murine leukemia virus reverse transcriptase and oligo(DT)₁₅ as a primer (Amersham Pharmacia, Piscataway, NJ).

PCR amplification and EST cloning

Primers used in this study are listed below in Table 6:

5

Table 6	
1. For 1st round PCR amplification	
Pem 5 (SEQ ID NO:25)	5'-GTGGACAAGAGGAAGCACAA-3' (derived from <i>OY-MS-4</i>)
EST-2 (SEQ ID NO:26)	5'-TCTCCCCATCTCACTCCAC-3' (derived from human testis EST clone, zt86b04.r1 [GenBank accession number AA397852])
2. For 2nd and 3rd PCR amplification, and analysis of mRNA expression in normal and malignant tissues	
ht-5 (SEQ ID NO:27)	5'-AAGGACAGGGGACTAAGGAG-3'
ht-3 (SEQ ID NO:28)	5'-CCGTACAAATCCAGCCCGTA-3'
ht-1 (SEQ ID NO:29)	5'-ATGTGAGTAGGGGCCGAGTA-3'
3. For 5' RACE	
GSP1-T1 (SEQ ID NO:30)	5'-TTCCTGGGCTGATCGAATGAG-3'
GSP2-T1 (SEQ ID NO:31)	5'-GCAAAAGAGGAAGGGTTAGAAG-3' (nested primer)
Tes-N1 (SEQ ID NO:32)	5'-CCGTGGTTTTTCATATTGGTC-3' (sequence primer)

A search was initiated for a human homologue gene of OY-MC-4 (*pem*), which had been found in murine sarcoma Meth A by SEREX analysis as described above. Human testis cDNA was amplified by PCR using primers pem5 (SEQ ID NO:25) from mouse OY-MC-4 and EST-2 (SEQ ID NO:26) from an EST database derived human testis clone. For first round PCR amplification, human testis cDNA was supplemented with 5 µl of 10x PCR buffer (100 mM Tris-HCl, 500 mM KCl, 0.01% (w/v) gelatin), 3 µl of 25 mM MgCl₂, 2 µl of 10 mM dNTPs, 5 µl of 5 µM each primer (pem5 and EST-2), 1.25 units of AmpliTaq DNA Polymerase (Perkin Elmer) and water to a final volume of 50 µl. PCR cycle conditions were 1 min at 94°C, 1 min at 54°C and 1.5 min at 72°C for 35 cycles. These cycles were preceded by 1 min of denaturation at 94°C and followed by a 10 min elongation step at 72°C.

A PCR product of 1.1 kb was obtained and sequenced using an ABI PRISM

automated sequencer. Homology walking was performed using the human dbEST database maintained by the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/dbEST>). Several ESTs corresponded to the PCR product. Most of the ESTs were cDNA clones derived from human testis. The EST clusters are shown in Fig. 1. The 3' end of the gene was extended by the sequence information of more than 11 ESTs which exhibited homology to the PCR product. Nucleotide sequence (nucleotides 261-1796 of SEQ ID NO:23) of the PCR product of 1.5 kb which was obtained using only anti-sense primer ht-1 (SEQ ID NO:29) was identical to the deduced sequence.

10 **5'-RACE amplification**

5' RACE was performed to identify the cDNA 5' end sequence using the 5' RACE System for Rapid Amplification kit following the manufacturer's protocols (Life Technologies, Inc., Rockville, MD). Total RNA from human testis was used as a template and first-strand cDNA was synthesized using specific primer, GSP1-T1 (SEQ ID NO:30).

15 The RACE product was sequenced using an ABI PRISM automated sequencer with sequence primer, Tes-N1 (SEQ ID NO:32).

Full length cDNA nucleotide sequence (SEQ ID NO:23) and deduced amino acid sequence (SEQ ID NO:24) are shown in Fig. 2.

20 **Homology**

The full length cDNA has homology with guinea pig and porcine acrosomal protein sp32 genes. Other homologous nucleic acid sequences are set forth in Table 8 below.

Expression profile of OY-TES-1

25 Specific primers, ht-5 (SEQ ID NO:27) and ht-3 (SEQ ID NO:28), were designed to amplify cDNA segments from normal tissue (MTC panels; Clontech, Palo Alto, CA) and tumor samples. RT-PCR was performed as described above for 30 amplification cycles with an annealing temperature of 62°C. Expression of the mRNA was observed with only testis in normal adult tissues and with various malignant tumors (Table 7).

Table 7: OY-TES-1 mRNA expression in human tumors

Tumor type	mRNA, positive/total
Bladder cancer	11/39
Lung cancer	4/11
Breast cancer	2/5
Hepatoma	2/5
Colon cancer	2/13
Stomach cancer	0/5
Renal cancer	0/10

These findings indicated that the cDNA identified is the human homologue of acrosomal protein sp32 and has the expression characteristics of a cancer/testis antigen.

5 Therefore, the cDNA was named OY-TES-1 antigen.

Table 8: Sequence homologies

10

SEQ ID NO:9

15

AL031055, U03904, AB021490.2, AF144629.1, U81385, X94191, X60820, M80939, M80938, W45845, AA276778, W12845, AA445681, AI174093, AA163016, AA791702, AA125123, AA793827, AA198900, AA003763, AI182954, AI597203.1, AA589600, AA561605, AV323570.1, AI614370.1, AA497714, AV233618.1, AW107276.1, AA754927, AA510149, W41547, AF039235, AA358918, AA373939, AI916936.1, AA315994, AA361102, AI440178.1, AI022281, AI743350.1, AA621367, AA598742, Z25343.

20

SEQ ID NO:13

25

NM_004592.1, J05080, AL034558.2, AC011001.2, AF168956.1, AB019001.1, AF126480.1, AC005142, AB015598.1, AF098161, AF101438.1, AF071506, AF071527.1, AB011167, U58748, L38477, Z11997, Y00365, U00431, L14429, L19249, AF003151.2, AP000346.1, AC005135, Z83122.1, L44140.1, X86810, D64111, L36829, K02587, NM_000117.1, X00654, X68946, AL132979.1, AC012380.1, AF109719.2, NM_006719.2, NM_002313.3, AB022692.1, NM_006720.1, AJ242956.1, NM_005691.1, NM_005378.1, AL034408.2, U71084, AJ131148.1, AF042714, AF076274, AC004005, Z93242, AF018165, AF025480, Z75710.1, AF005370, AF005654, Z81515.1, Z78416.1, Z72686, Z71584, X59601, L38856, U33758, D31967, Z48618, Z46259, X59863, X57561, X17199, Z21611, X07699, D26114, X03295, X02363, Y00664, Z22572, NM_004606.1, M32092, M13241, L23114, M73444, L19597, L27631, L09209, D31883, NM_001642.1, X74225.1, M22580, M23216, M34271, M13228, D21877, H44689, AA027262, AA256873, AI871457.1, AI950925.1, AA812028, AI431592.1, AA374595, AA437048, AA282462, N84830, R70432, AI417091.1, AI391464, AI016313, AA548044, AA484436, AA442000, W52798, W25804, W24038, N49505, AW005182.1, AW002246.1,

35

AI937788.1, AI937103.1, AL044552.1, AL041478.1, AI696132.1, AL047599.1,
AI499031.1, AI371599, AI368947, AI368871, AI337311, AI334569, AI255139,
AI242685, AA725144, AA693685, AA663606, AA578529, AA315123, AA203519,
AA167831, AA083111, AA057640, AA203468, W67602, W67601, N95426, R80498,
5 R79280, R66447, R52824, R33900, AI316495, AA420323, AI173226, AW045177.1,
C80075, AW049212.1, AW045819.1, AW048739.1, C76919, AA254996, C87870,
AA030824, AA561021, AI573843.1, AA030365, W08619, AA416313, AV330281.1,
AA204230, AV277735.1, AV094608.1, AI152993, AV031612.1, AV254814.1,
AV276238.1, AV315661.1, AV274035.1, AV240388.1, AV365468.1, AV374341.1,
10 AV298568.1, AV330260.1, AV305225.1, AI154346, AV225442.1, AI839392.1,
AV326271.1, AV349170.1, AV247245.1, AA816058, AV257049.1, AV235616.1,
AV322742.1, AV080798.1, AA068921, AV226134.1, AV275975.1, AV248838.1,
W82727, AA217762, D18136, AA789824, AI527541, AI227319, W64508, AA797091,
AA239762, AA823138, AA475558, AA197732, W70850, AI956759.1, AV170718.1,
15 AI791067.1, AI790663.1, AV125900.1, AV122779.1, AV096937.1, AI648759.1,
AI550256.1, AI463840, AU015737, AU015657, AU014623, AA895398, AA798375,
AA797955, AA793428, C81382, C80585, C80564, AA607305, AA588982, AA571572,
AA560832, AA474262, AA467444, AA413093, AA275567, AA123213, AA120232,
AA111692, W97542, AI842860.1, AA268371, AA031077, AV294981.1, AA833307,
20 AA472076, AI711096.1, AI709709.1, AI233561, AI180068, AI170909, AI102461,
AI576455.1, AA957106.1, AI008470, AA944954, AI043740, AI175545, H33858,
AW165033.1, AW128282.1, AW019641.1, AI779631.1, AI774248.1, AI773240.1,
AW165272.1, AW096676.1, AW091803.1, AI722622.1, AI722217.1, AI722132.1,
AI721578.1, AU062029.1, AI728625.1, AI542408, AI516896, AA605835, C32241.1,
25 D67546, AV384376.1, AV383428.1, AI947408.1, AI776807.1, AI626286.1,
AI621775.1, AI586631.1, AI545013, C91835, AF020720, AA494586, T88263,
T13720, T04634, R02924.

30 SEQ ID NO:15

NM_005003.1, X61997, AC002400, AF131865.1, AC002526, D17577, AL023755.5,
AC006972.2, AL034489, Y16789, AC004512, Z74979, L40557, AI648903.1,
AI119292, AW107104.1, AI647845.1, AI119359, AI929857.1, AI849803.1,
35 AI047860, AI850286.1, AV000514.1, AI046666, AI413300, W89503, AA939621,
AA210326, AA168999, AA185613, AA929298, AA212234, AI152408, AA915295,
AI172915, AA790460, AI930104.1, AA466529, AA032743, AA048160, AA711023,
AV006031.1, AA738875, AA212314, AA197653, AA030594, W56963, AI227424,
AI893356.1, AA271005, AA060580, W42142, AA119387, W79962, AI115431,
40 AA822668, AA007973, W80240, W13086, W12715, AI877179.1, AA013975, W15883,
AA500928, AA238199, W85108, AA930915, AA240213, W40680, AI117345, W76948,
W50674, AI508285, W82342, AA110994, W43957, AA646522, AA437633, AA989964,
AA178192, AA920691, W48200, AV000328.1, AA271056, AV066478.1, D17880,
AA684079, W83559, AV021796.1, AA204050, AA062534, AA033484, AA823129,
45 AA162638, AV060641.1, AV083684.1, W16252, AV118462.1, R75584, AV091441.1,
AV080619.1, AV056483.1, AV051905.1, AV085492.1, AA207625, AV085512.1,
AV033153.1, AV006258.1, AV062958.1, AV084081.1, AV225927.1, AV152111.1,
AA690970, AA075950, AA082003, AA074065, AA313368, AW007505.1, AI889052.1,
AI598144.1, AI095590, AI017929, AA310175, AA873566, AA868795, AA868719,
50 AA777631, AA600754, AA476531, AW044220.1, F36770.1, F25329.1, AI083718,
AA993399, AA935632, AA618021, AA459170, N42432, N33459, AI873496.1,
F24032.1, AI581853.1, AI302612, AI301635, AI285263, AI248971, AI220450,
AI160021, AI128630, AI032546, AI016702, AI014825, AA961682, AA936142,
AA860514, AA833995, AA831482, AA447569, AA417135, AW166336.1, AW073092.1,
55 AI863340.1, AI520938, AA829435, AA508601, AI815674.1, AI720708.1, AI161342,
AA875886, AA983159, AA587029, AA689457, AI469794.1, AA729752, AA448553,
AA909264, AA868027, AA507444, AA416980, AA886275, AA883939, AA878640,
AI469742, AI281705, AA885866, AW139387.1, AA152181, AA729741, N35532,
AA723449, N94436, AA936986, AA917976, AA632370, AA150058, N35132, N26481,
60 AA909159, AA639717, AA642285, N35190, AA906828, AI609646.1, AA827419,
AA689510, AA872218, AA136946, F18960.2, N86284, D54530, AI352654, AA152029,
AI525049.1.

65 SEQ ID NO:17

U06944, U88344, AC005261, AC004073, AC005966, U34740, AF056562.1,

AF056561.1, AB026654.1, AL034418.2, AJ011712.1, AC005546, U92322,
AL022721.1, AL021878, AC003058, Z99102, AB003505, AB003504, AE001099,
AB000493, U49391, U60211, Z69902.1, M34652, AI596060.1, AA272122, AA269508,
AI849761.1, AI874995.1, AU024159, AA270426, AU079534.1, AA475684, W15945,
5 AA590308, AI956733.1, AA518797, AI847841.1, AI852328.1, AA119752,
AV204180.1, AV306904.1, W87234, AV355049.1, AV320983.1, AI882238.1,
AV299331.1, AV033250.1, AV032030.1, AV269232.1, AA170872, AV301048.1,
AV226430.1, AV355108.1, AI663065.1, AV295507.1, AA119300, AA116521,
AI882051.1, AV364043.1, AV224385.1, AV337033.1, AV248698.1, AW123672.1,
10 AV282468.1, AV214423.1, AV309192.1, AV151663.1, AV125955.1, AV106869.1,
AV098772.1, AV077295.1, AV054772.1, AV027590.1, AV014774.1, AV013646.1,
AV214005.1, AV066448.1, AV026120.1, AV009189.1, AV336813.1, AV335057.1,
AV331939.1, AV269945.1, AV227030.1, AV161933.1, AV130365.1, AV125700.1,
AV121841.1, AV101405.1, AV096440.1, AV089404.1, AV076355.1, AV069807.1,
15 AV066548.1, AV064183.1, AV061514.1, AV055563.1, AV051613.1, AV006337.1,
AI481849, AI948435.1, AI912734.1, AI796967.1, AI740869.1, AI498701.1,
AI197956, AI197904, AI003139, AA887501, AA191594, W86389, N64011, H99275,
AI985257.1, AA346169, AA219621, AI282275, AA191083, AA779217, D61002,
H05031, H81379, R20553, R41899, AI864073.1, AA992514, D29534, W92008,
20 AA393678, W92009.

SEQ ID NO:23

25 D16200.1, AF085884.1, D16203.1, G23540.1, D17574.1, D17573.1, AC018363.6,
AF113672.1, AL117419.1, Y15994.1, AF148693.1, AE003584.1, AC002460.1,
AC004716.1, AL033504.3, X52027.1, D85605.1, D86959.1, AB002804.1,
AA864327.1, AA824340.1, AA496077.1, AI125678.1, AI028208.1, AI126598.1,
AA406076.1, AI769240.1, AI575049.1, AI333870.1, AI333624.1, AI141116.1,
30 AI131223.1, AI091519.1, AI122673.1, AA709069.1, AA443593.1, AI147148.1,
AA471063.1, H30251.1, R07741.1, AI138894.1, AI024421.1, AA993247.1,
AW592506.1, AA424694.1, AA405433.1, AA473853.1, H30315.1, AI382680.1,
AI574731.1, R07740.1, AA992230.1, AI326707.1, AA154158.1, AI125674.1,
AW358264.1, AI573445.1, AL042116.1, AV005781.1, AI047576.1, AI553526.1,
35 AA104771.1, AA065355.1, AI208979.1, AI716721.1, AA621303.1, AV040492.2.

EQUIVALENTS

Those skilled in the art will recognize, or be able to ascertain using no more than
40 routine experimentation, many equivalents to the specific embodiments of the invention
described herein. Such equivalents are intended to be encompassed by the following claims.

All references disclosed herein are incorporated by reference in their entirety.

We claim:

Claims

1. A method of diagnosing a disorder characterized by expression of a cancer associated antigen precursor coded for by a nucleic acid molecule, comprising:

contacting a biological sample isolated from a subject with an agent that specifically
5 binds to the nucleic acid molecule, an expression product thereof, or a fragment of an expression product thereof complexed with a MHC molecule, wherein the nucleic acid molecule is a NA Group 1 molecule, and

determining the interaction between the agent and the nucleic acid molecule or the expression product as a determination of the disorder.

10

2. The method of claim 1, wherein the agent is selected from the group consisting of

(a) a nucleic acid molecule comprising NA group 1 molecules or a fragment thereof,

(b) a nucleic acid molecule comprising NA group 3 molecules or a fragment
15 thereof,

(c) a nucleic acid molecule comprising NA group 5 molecules or a fragment thereof,

(d) an antibody that binds to an expression product of NA group 1 molecules,

(e) an antibody that binds to an expression product of NA group 3 molecules,

20 (f) an antibody that binds to an expression product of NA group 5 molecules,

(g) an agent that binds to a complex of a MHC molecule and a fragment of an expression product of a NA group 1 molecule,

(h) an agent that binds to a complex of a MHC molecule and a fragment of an expression product of a NA group 3 molecule, and

25 (i) an agent that binds to a complex of a MHC molecule and a fragment of an expression product of a NA group 5 molecule.

3. The method of claim 1, wherein the disorder is characterized by expression of a plurality of cancer associated antigen precursors and wherein the agent is a plurality of agents,
30 each of which is specific for a different cancer associated antigen precursor, and wherein said plurality of agents is at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, or at least 8, at least 9 or at least 10 such agents.

4. The method of claims 1-3, wherein the agent is specific for a cancer associated antigen precursor that is a fibrosarcoma cancer associated antigen precursor.

5. The method of claim 2, wherein the NA group 1 molecule is SEQ ID NO:23.

5

6. A method for determining regression, progression or onset of a condition characterized by expression of abnormal levels of a protein encoded by a nucleic acid molecule that is a NA Group 1 molecule, comprising

monitoring a sample, from a patient who has or is suspected of having the condition,
10 for a parameter selected from the group consisting of

- (i) the protein,
- (ii) a peptide derived from the protein,
- (iii) an antibody which selectively binds the protein or peptide, and
- (iv) cytolytic T cells specific for a complex of the peptide derived from the

15 protein and an MHC molecule,
as a determination of regression, progression or onset of said condition.

7. The method of claim 6, wherein the sample is a body fluid, a body effusion or a tissue.

20 8. The method of claim 6, wherein the step of monitoring comprises contacting the sample with a detectable agent selected from the group consisting of

- (a) an antibody which selectively binds the protein of (i), or the peptide of (ii),
- (b) a protein or peptide which binds the antibody of (iii), and
- (c) a cell which presents the complex of the peptide and MHC molecule of (iv).

25

9. The method of claim 8, wherein the antibody, the protein, the peptide or the cell is labeled with a radioactive label or an enzyme.

10. The method of claim 6, comprising assaying the sample for the peptide.

30

11. The method of claim 6, wherein the nucleic acid molecule is a NA Group 3 molecule.

12. The method of claim 6, wherein the nucleic acid molecule is a NA Group 5 molecule.

13. The method of claim 6, wherein the protein is a plurality of proteins, the parameter is a plurality of parameters, each of the plurality of parameters being specific for a different of the plurality of proteins, at least one of which is a cancer associated protein encoded by a NA
5 Group 1 molecule.
14. The method of claim 6, wherein the NA Group 1 molecule is SEQ ID NO:23.
15. A pharmaceutical preparation comprising
10 an agent which when administered to the subject enriches selectively the presence of complexes of a MHC molecule and a cancer associated antigen, and
a pharmaceutically acceptable carrier, wherein the cancer associated antigen is a fragment of a cancer associated antigen precursor encoded by a nucleic acid molecule comprising a NA Group 1 molecule.
15
16. The pharmaceutical preparation of claim 15, wherein the agent comprises a plurality of agents, each of which enriches selectively in the subject complexes of a MHC molecule and a different cancer associated antigen, wherein at least one of the cancer associated antigens is encoded by a NA Group 1 molecule.
20
17. The pharmaceutical preparation of claim 16, wherein the plurality is at least two, at least three, at least four or at least 5 different such agents.
18. The pharmaceutical preparation of claim 15, wherein the nucleic acid molecule is a
25 NA Group 3 nucleic acid molecule.
19. The pharmaceutical preparation of claim 15, wherein the agent is selected from the group consisting of
(1) an isolated polypeptide comprising the cancer associated antigen, or a functional
30 variant thereof,
(2) an isolated nucleic acid operably linked to a promoter for expressing the isolated polypeptide, or functional variant thereof,
(3) a host cell expressing the isolated polypeptide, or functional variant thereof, and

(4) isolated complexes of the polypeptide, or functional variant thereof, and a MHC molecule.

20. The pharmaceutical preparation of claim 15, wherein the NA Group 1 molecule is
5 SEQ ID NO:23.

21. The pharmaceutical preparation of claims 15-20, further comprising an adjuvant.

22. The pharmaceutical preparation of claim 15, wherein the agent is a cell expressing an
10 isolated polypeptide comprising the cancer associated antigen or a functional variant thereof,
and wherein the cell is nonproliferative.

23. The pharmaceutical preparation of claim 15, wherein the agent is a cell expressing an
isolated polypeptide comprising the cancer associated antigen or a functional variant thereof,
15 and wherein the cell expresses a MHC molecule that binds the polypeptide.

24. The pharmaceutical preparation of claim 15, wherein the agent is at least two, at least
three, at least four or at least five different polypeptides, each coding for a different cancer
associated antigen or functional variant thereof, wherein at least one of the cancer associated
20 antigens is encoded by a NA Group 1 molecule.

25. The pharmaceutical preparation of claim 15, wherein the agent is a PP Group 2
polypeptide.

25 26. The pharmaceutical preparation of claim 15, wherein the agent is a PP Group 3
polypeptide or a PP Group 4 polypeptide.

27. The pharmaceutical preparation of claim 23, wherein the cell expresses one or both of
the polypeptide and the MHC molecule recombinantly.

30

28. The pharmaceutical preparation of claim 23, wherein the cell is nonproliferative.

29. A composition comprising

an isolated agent that binds selectively a PP Group 1 polypeptide.

30. The composition of claim 29, wherein the PP Group 1 polypeptide is SEQ ID NO:24.

5 31. The composition of matter of claim 26, wherein the agent binds selectively a PP Group 2 polypeptide.

32. The composition of matter of claim 29, wherein the agent binds selectively a PP Group 3 polypeptide.

10

33. The composition of matter of claim 29, wherein the agent binds selectively a PP Group 4 polypeptide.

15

34. The composition of matter of claim 29, wherein the agent binds selectively a PP Group 5 polypeptide.

35. The composition of claims 29-34, wherein the agent is a plurality of different agents that bind selectively at least two, at least three, at least four, or at least five different such polypeptides.

20

36. The composition of claims 29-34, wherein the agent is an antibody.

37. The composition of claim 35, wherein the agent is an antibody.

25

38. A composition of matter comprising
a conjugate of the agent of claims 29-34 and a therapeutic or diagnostic agent.

39. A composition of matter comprising
a conjugate of the agent of claim 35 and a therapeutic or diagnostic agent.

30

40. The composition of matter of claim 38, wherein the conjugate is of the agent and a therapeutic or diagnostic that is a toxin.

US 2016/0265561 A1

41. A pharmaceutical composition comprising an isolated nucleic acid molecule selected from the group consisting of NA Group 1 molecules and NA Group 2 molecules, and a pharmaceutically acceptable carrier.

5 42. The pharmaceutical composition of claim 41, wherein the NA Group 1 molecule is SEQ ID NO:23.

43. The pharmaceutical composition of claim 41, wherein the isolated nucleic acid molecule comprises a NA Group 3 or NA Group 4 molecule.

10

44. The pharmaceutical composition of claim 41, wherein the isolated nucleic acid molecule comprises at least two isolated nucleic acid molecules coding for two different polypeptides, each polypeptide comprising a different cancer associated antigen.

15 45. The pharmaceutical composition of claims 41-44 further comprising an expression vector with a promoter operably linked to the isolated nucleic acid molecule.

46. The pharmaceutical composition of claims 41-44 further comprising a host cell recombinantly expressing the isolated nucleic acid molecule.

20

47. A pharmaceutical composition comprising
an isolated polypeptide comprising a PP Group 1 or a PP Group 2 polypeptide, and
a pharmaceutically acceptable carrier.

25 48. The pharmaceutical composition of claim 47, wherein the PP Group 1 polypeptide is SEQ ID NO:24.

49. The pharmaceutical composition of claim 47, wherein the isolated polypeptide comprises a PP Group 3 or a PP Group 4 polypeptide.

30

50. The pharmaceutical composition of claim 47, wherein the isolated polypeptide comprises at least two different polypeptides, each comprising a different cancer associated antigen.

51. The pharmaceutical composition of claim 47, wherein the isolated polypeptides are PP Group 3 polypeptides or MHC binding fragments thereof.

5 52. The pharmaceutical composition of claim 47, wherein the isolated polypeptides are PP Group 5 polypeptides or MHC binding fragments thereof.

53. The pharmaceutical composition of claims 47-52, further comprising an adjuvant.

10 54. An isolated nucleic acid molecule comprising a NA Group 3 molecule.

55. An isolated nucleic acid molecule comprising a NA Group 4 molecule.

56. An isolated nucleic acid molecule selected from the group consisting of

15 (a) a fragment of a nucleic acid molecule having a nucleotide sequence selected from the group consisting of nucleotide sequences set forth as SEQ ID NOs. 9, 13, 15, 17, 19, and 23, of sufficient length to represent a sequence unique within the mouse or human genomes, and identifying a nucleic acid encoding a cancer associated antigen precursor,

(b) complements of (a),

20 provided that the fragment includes a sequence of contiguous nucleotides which is not identical to any sequence selected from the sequence group consisting of

(1) sequences having the GenBank accession numbers of Table 8,

(2) complements of (1), and

(3) fragments of (1) and (2).

25 57. The isolated nucleic acid molecule of claim 56, wherein the sequence of contiguous nucleotides is selected from the group consisting of:

(1) at least two contiguous nucleotides nonidentical to the sequence group,

(2) at least three contiguous nucleotides nonidentical to the sequence group,

30 (3) at least four contiguous nucleotides nonidentical to the sequence group,

(4) at least five contiguous nucleotides nonidentical to the sequence group,

(5) at least six contiguous nucleotides nonidentical to the sequence group,

(6) at least seven contiguous nucleotides nonidentical to the sequence group.

58. The isolated nucleic acid molecule of claim 56, wherein the fragment has a size selected from the group consisting of at least: 8 nucleotides, 10 nucleotides, 12 nucleotides, 14 nucleotides, 16 nucleotides, 18 nucleotides, 20, nucleotides, 22 nucleotides, 24
5 nucleotides, 26 nucleotides, 28 nucleotides, 30 nucleotides, 50 nucleotides, 75 nucleotides, 100 nucleotides, and 200 nucleotides.
59. The isolated nucleic acid molecule of claim 50, wherein the molecule encodes a polypeptide which, or a fragment of which, binds a MHC receptor or an antibody.
- 10 60. An expression vector comprising an isolated nucleic acid molecule of any of claims 54-59 operably linked to a promoter.
61. An expression vector comprising a nucleic acid operably linked to a promoter, wherein
15 the nucleic acid is a NA Group 2 molecule.
62. An expression vector comprising a NA Group 1 or Group 2 molecule and a nucleic acid encoding a MHC molecule.
- 20 63. The expression vector of claim 62, wherein the NA Group 1 molecule is SEQ ID NO:23.
64. A host cell transformed or transfected with an expression vector of claim 60.
- 25 65. A host cell transformed or transfected with an expression vector of any of claims 61-63.
66. A host cell transformed or transfected with an expression vector of claim 60 and further comprising a nucleic acid encoding a MHC molecule.
- 30 67. A host cell transformed or transfected with an expression vector of claim 61 and further comprising a nucleic acid encoding a MHC molecule.

68. An isolated polypeptide encoded by the isolated nucleic acid molecule of claim 54 or claim 55.

69. A fragment of the polypeptide of claim 68 which is immunogenic.

5

70. The fragment of claim 69, wherein the fragment, or a portion of the fragment, binds a MHC molecule or an antibody.

71. An isolated fragment of a cancer associated antigen precursor which, or portion of which, binds a MHC molecule or an antibody, wherein the precursor is encoded by a nucleic acid molecule that is a NA Group 1 molecule.

10

72. The isolated fragment of claim 71, wherein the NA Group 1 molecule is SEQ ID NO:23.

15

73. The fragment of claim 71, wherein the fragment is part of a complex with a MHC molecule.

20

74. The fragment of claim 73, wherein the fragment is between 8 and 12 amino acids in length.

75. An isolated polypeptide comprising a fragment of the polypeptide of claim 68 of sufficient length to represent a sequence unique within the mouse or human genomes and identifying a polypeptide that is a cancer associated antigen precursor.

25

76. A kit for detecting the presence of the expression of a cancer associated antigen precursor comprising

30

a pair of isolated nucleic acid molecules each of which consists essentially of a molecule selected from the group consisting of (a) a 12-32 nucleotide contiguous segment of the nucleotide sequence of any of the NA Group 1 molecules and (b) complements of (a), wherein the contiguous segments are nonoverlapping.

77. The kit of claim 76, wherein the NA Group 1 molecule is SEQ ID NO:23.

78. The kit of claim 76, wherein the pair of isolated nucleic acid molecules is constructed and arranged to selectively amplify an isolated nucleic acid molecule that is a NA Group 3 molecule.

5

79. A method for treating a subject with a disorder characterized by expression of a cancer associated antigen precursor, comprising

administering to the subject an amount of an agent, which enriches selectively in the subject the presence of complexes of a MHC molecule and a cancer associated antigen, effective to ameliorate the disorder, wherein the cancer associated antigen is a fragment of a cancer associated antigen precursor encoded by a nucleic acid molecule selected from the group consisting of

10

- (a) a nucleic acid molecule comprising NA Group 1 molecules,
- (b) a nucleic acid molecule comprising NA Group 3 molecules,
- (c) a nucleic acid molecule comprising NA Group 5 molecules.

15

80. The method of claim 79, wherein the NA Group 1 molecule is SEQ ID NO:23.

81. The method of claim 79, wherein the disorder is characterized by expression of a plurality of cancer associated antigen precursors and wherein the agent is a plurality of agents, each of which enriches selectively in the subject the presence of complexes of a MHC molecule and a different cancer associated antigen, wherein at least one of the cancer associated antigens is encoded by a NA Group 1 molecule.

20

82. The method of claim 81, wherein the plurality is at least 2, at least 3, at least 4, or at least 5 such agents.

25

83. The method of claims 79-82, wherein the agent is an isolated polypeptide selected from the group consisting of PP Group 1, PP Group 2, PP Group 3, PP Group 4 and PP Group 5.

30

84. The method of claims 79-82, wherein the disorder is cancer.

85. The method of claims 83, wherein the disorder is cancer.

86. A method for treating a subject having a condition characterized by expression of a cancer associated antigen precursor in cells of the subject, comprising:

- 5 (i) removing an immunoreactive cell containing sample from the subject,
- (ii) contacting the immunoreactive cell containing sample to the host cell under conditions favoring production of cytolytic T cells against a cancer associated antigen which is a fragment of the precursor,
- (iii) introducing the cytolytic T cells to the subject in an amount effective to lyse
- 10 cells which express the human cancer associated antigen, wherein the host cell is transformed or transfected with an expression vector comprising an isolated nucleic acid molecule operably linked to a promoter, the isolated nucleic acid molecule being selected from the group of nucleic acid molecules consisting of NA Group 1 molecules, NA Group 2 molecules, NA Group 3 molecules, NA Group 4 molecules, and NA Group 5 molecules.

15 87. The method of claim 86, wherein the NA Group 1 molecule is SEQ ID NO:23.

88. The method of claim 86, wherein the host cell recombinantly expresses a MHC molecule which binds the cancer associated antigen.

20 89. The method of claim 86, wherein the host cell endogenously expresses a MHC molecule which binds the cancer associated antigen.

90. A method for treating a subject having a condition characterized by expression of a cancer associated antigen precursor in cells of the subject, comprising:

- 25 (i) identifying a nucleic acid molecule expressed by the cells associated with said condition, wherein said nucleic acid molecule is a NA Group 1 molecule;
- (ii) transfecting a host cell with a nucleic acid selected from the group consisting of (a) the nucleic acid molecule identified, (b) a fragment of the nucleic acid identified which
- 30 includes a segment coding for a cancer associated antigen, (c) deletions, substitutions or additions to (a) or (b), and (d) degenerates of (a), (b), or (c);
- (iii) culturing said transfected host cells to express the transfected nucleic acid molecule, and;

(iv) introducing an amount of said host cells or an extract thereof to the subject effective to increase an immune response against the cells of the subject associated with the condition.

5 91. The method of claim 90, wherein the NA Group 1 molecule is SEQ ID NO:23.

92. The method of claim 90, further comprising identifying a MHC molecule which presents a portion of an expression product of the nucleic acid molecule, wherein the host cell expresses the same MHC molecule as identified and wherein the host cell presents a MHC
10 binding portion of the expression product of the nucleic acid molecule.

93. The method of claim 90, wherein the immune response comprises a B-cell response or a T-cell response.

15 94. The method of claim 93, wherein the response is a T-cell response which comprises generation of cytolytic T-cells specific for the host cells presenting the portion of the expression product of the nucleic acid molecule or cells of the subject expressing the cancer associated antigen.

20 95. The method of claim 90, wherein the nucleic acid molecule is a NA Group 3 molecule.

96. The method of any of claims 90-92, further comprising treating the host cells to render them non-proliferative.

25 97. A method for treating or diagnosing or monitoring a subject having a condition characterized by expression of an abnormal amount of a protein encoded by a nucleic acid molecule that is a NA Group 1 molecule, comprising

administering to the subject an antibody which specifically binds to the protein or a peptide derived therefrom, the antibody being coupled to a therapeutically useful agent, in an
30 amount effective to treat the condition.

98. The method of claim 97, wherein the NA Group 1 molecule is SEQ ID NO:23.

99. The method of claim 97, wherein the antibody is a monoclonal antibody.

100. The method of claim 99, wherein the monoclonal antibody is a chimeric antibody or a humanized antibody.

5

101. A method for treating a condition characterized by expression in a subject of abnormal amounts of a protein encoded by a nucleic acid molecule that is a NA Group 1 nucleic acid molecule, comprising

administering to a subject a pharmaceutical composition of any one of claims 13-25
10 and 37-47 in an amount effective to prevent, delay the onset of, or inhibit the condition in the subject.

102. The method of claim 101, wherein the condition is cancer.

15 103. The method of claim 101, further comprising first identifying that the subject expresses in a tissue abnormal amounts of the protein.

104. The method of claim 102, further comprising first identifying that the subject expresses in a tissue abnormal amounts of the protein.

20

105. A method for treating a subject having a condition characterized by expression of abnormal amounts of a protein encoded by a nucleic acid molecule that is a NA Group 1 nucleic acid molecule, comprising

(i) identifying cells from the subject which express abnormal amounts of the protein;
25 (ii) isolating a sample of the cells;
(iii) cultivating the cells, and
(iv) introducing the cells to the subject in an amount effective to provoke an immune response against the cells.

30 106. The method of claim 105, wherein the NA Group 1 molecule is SEQ ID NO:23.

107. The method of claim 105, further comprising rendering the cells non-proliferative, prior to introducing them to the subject.

108. A method for treating a pathological cell condition characterized by aberrant expression of a protein encoded by a nucleic acid molecule that is a NA Group 1 nucleic acid molecule, comprising

5 administering to a subject in need thereof an effective amount of an agent which inhibits the expression or activity of the protein.

109. The method of claim 108, wherein the NA Group 1 molecule is SEQ ID NO:23.

10 110. The method of claim 108, wherein the agent is an inhibiting antibody which selectively binds to the protein and wherein the antibody is a monoclonal antibody, a chimeric antibody or a humanized antibody.

111. The method of claim 108, wherein the agent is an antisense nucleic acid molecule
15 which selectively binds to the nucleic acid molecule which encodes the protein.

112. The method of claim 108, wherein the nucleic acid molecule is a NA Group 3 nucleic acid molecule.

20 113. A composition of matter useful in stimulating an immune response to a plurality of a proteins encoded by nucleic acid molecules that are NA Group 1 molecules, comprising a plurality of peptides derived from the amino acid sequences of the proteins, wherein the peptides bind to one or more MHC molecules presented on the surface of the cells which express an abnormal amount of the protein.

25 114. The composition of matter of claim 113, wherein the NA Group 1 molecule is SEQ ID NO:23.

115. The composition of matter of claim 113, wherein at least a portion of the plurality of
30 peptides bind to MHC molecules and elicit a cytolytic response thereto.

116. The composition of matter of claim 115, further comprising an adjuvant.

117. The composition of matter of claim 116, wherein said adjuvant is a saponin, GM-CSF, or an interleukin.

118. The composition of matter of claim 113, further comprising at least one peptide useful
5 in stimulating an immune response to at least one protein which is not encoded by nucleic acid molecules that are NA Group 1 molecules, wherein the at least one peptide binds to one or more MHC molecules.

119. An isolated antibody which selectively binds to a complex of:

10 (i) a peptide derived from a protein encoded by a nucleic acid molecule that is a NA Group 1 molecule and

(ii) and an MHC molecule to which binds the peptide to form the complex,
wherein the isolated antibody does not bind to (i) or (ii) alone.

120. The isolated antibody of claim 119, wherein the NA Group 1 molecule is SEQ ID
15 NO:23.

121. The antibody of claim 119, wherein the antibody is a monoclonal antibody, a chimeric
antibody, a humanized antibody, or a fragment thereof.

20

Abstract

Cancer associated antigens have been identified by autologous antibody screening of libraries of nucleic acids expressed in methylcholanthrene-induced fibrosarcoma cancer cells
5 using antisera from mice bearing such tumors. The invention relates to nucleic acids and encoded polypeptides which are cancer associated antigens expressed in mice afflicted with methylcholanthrene-induced fibrosarcomas, as well as homologs thereof, particularly human homologs. The invention provides, *inter alia*, isolated nucleic acid molecules, expression vectors containing those molecules and host cells transfected with those molecules. The
10 invention also provides isolated proteins and peptides, antibodies to those proteins and peptides and cytotoxic T lymphocytes which recognize the proteins and peptides. Fragments of the foregoing including functional fragments and variants also are provided. Kits containing the foregoing molecules additionally are provided. The molecules provided by the invention can be used in the diagnosis, monitoring, research, or treatment of conditions
15 characterized by the expression of one or more cancer associated antigens.

SEQUENCE LISTING

5 <110> Ono, Toshiro
Nakayama, Eiichi

<120> CANCER ASSOCIATED ANTIGENS AND USES
THEREFOR

10 <130> L0461/7086

<150> US 60/168,353
<151> 1999-12-01

15 <160> 32

<170> FastSEQ for Windows Version 3.0

20 <210> 1
<211> 538
<212> DNA
<213> Mus musculus

<400> 1

25	agatcaaggg	gaaaaggaga	accccatgcg	ggaactgcgc	atccgcaagc	tctgcctcaa	60
	tatctgcgtc	ggggagagcg	gagacagact	gacccgggca	gccaaggtgt	tggagcagct	120
	cacaggccag	accccggtgt	tctccaaagc	tagatacact	gtcaggtoct	ttggcatccg	180
	gagaaatgag	aagattgctg	ttcactgcac	agtccgcgga	gccaaggcag	aggaaattct	240
	ggagaaaggc	ctgaaggtgc	gggagtatga	gttgcggaaa	aataacttct	cggatactgg	300
	aaacttttgt	tttgggaattc	aagaacacat	tgacctgggc	atcaaatacg	acccaagcat	360
30	tgggatctac	ggcctggact	tctatgtggt	gctgggtagg	ccagggttca	gcacgcgaga	420
	caagaagcgc	agaacaggct	gcattggggc	caaacacaga	atcagcaagg	aggaggccat	480
	gcgctgggtc	cagcagaagt	acgatggaat	catccttctc	ggaaaataaa	cttgatcc	538

<210> 2

35 <211> 175
<212> PRT
<213> Mus musculus

<400> 2

40	Asp	Gln	Gly	Glu	Lys	Glu	Asn	Pro	Met	Arg	Glu	Leu	Arg	Ile	Arg	Lys
	1				5					10				15		
	Leu	Cys	Leu	Asn	Ile	Cys	Val	Gly	Glu	Ser	Gly	Asp	Arg	Leu	Thr	Arg
			20						25				30			
45	Ala	Ala	Lys	Val	Leu	Glu	Gln	Leu	Thr	Gly	Gln	Thr	Pro	Val	Phe	Ser
			35					40				45				
	Lys	Ala	Arg	Tyr	Thr	Val	Arg	Ser	Phe	Gly	Ile	Arg	Arg	Asn	Glu	Lys
	50						55			60						
	Ile	Ala	Val	His	Cys	Thr	Val	Arg	Gly	Ala	Lys	Ala	Glu	Glu	Ile	Leu
	65				70				75					80		
50	Glu	Lys	Gly	Leu	Lys	Val	Arg	Glu	Tyr	Glu	Leu	Arg	Lys	Asn	Asn	Phe
				85					90					95		
	Ser	Asp	Thr	Gly	Asn	Phe	Gly	Phe	Gly	Ile	Gln	Glu	His	Ile	Asp	Leu
			100				105						110			
	Gly	Ile	Lys	Tyr	Asp	Pro	Ser	Ile	Gly	Ile	Tyr	Gly	Leu	Asp	Phe	Tyr
55			115				120					125				
	Val	Val	Leu	Gly	Arg	Pro	Gly	Phe	Ser	Ile	Ala	Asp	Lys	Lys	Arg	Arg
			130				135				140					
	Thr	Gly	Cys	Ile	Gly	Ala	Lys	His	Arg	Ile	Ser	Lys	Glu	Glu	Ala	Met
	145				150				155					160		
60	Arg	Trp	Phe	Gln	Gln	Lys	Tyr	Asp	Gly	Ile	Ile	Leu	Pro	Gly	Lys	
				165					170					175		

<210> 3
<211> 1228
<212> DNA
<213> Mus musculus

5

<400> 3

	acagccgcat	cttcttgtgc	agtgccagcc	tcgtcccgtg	gacaaaatgg	tgaaggctcg	60
	tgtgaacgga	tttgccgta	ttgggcgcct	ggtcaccagg	gctgccattt	gcagtgccaa	120
	agtgagatt	gttgccatca	acgaccctt	cattgacctc	aactacatgg	tctacatgtt	180
10	ccagtatgac	tccactcacg	gcaaattcaa	cggcacagtc	aaggccgaga	atgggaagct	240
	tgtcatcaac	gggaagccca	tcaccatctt	ccaggagcga	gaccccaacta	acatcaaattg	300
	gggtgaggcc	ggtgctgagt	atgtcgtgga	gtctactggg	gtcttcacca	ccatggagaa	360
	ggccggggcc	cacttgaagg	gtggagccaa	acgggtcatc	atctccgccc	cttctgccga	420
	tgcccccatg	tttgtgatgg	gtgtgaacca	cgagaaatat	gacaactcac	tcaagattgt	480
15	cagcaatgca	tcttgacca	ccaactgctt	agccccctg	gccaaaggta	tccatgacaa	540
	ctttggcatt	gtggaaggcc	tcatgaccac	agtccatgcc	atcactgcca	cccagaagac	600
	tgtggatggc	ccctctggaa	agctgtggcg	tgatggccgt	ggggctgccc	agaacatcat	660
	ccctgcatcc	actggtgctg	ccaaggctgt	gggcaaggtc	atcccagagc	tgaacgggaa	720
	gctcactggc	atggccttcc	gtgttcttac	ccccaatgtg	tccgtcgtgg	atctgacgtg	780
20	ccgcctggag	aaacctgcca	agtatgatga	catcaagaag	gtggtgaagc	aggcatctga	840
	gggcccactg	aagggcattc	tgggctacac	tgaggaccag	gttgtctcct	gcgacttcaa	900
	cagcaactcc	cactcttcca	ccttcgatgc	cggggctggc	attgctctca	atgacaactt	960
	tgtcaagctc	atttctctgt	atgacaatga	atacggctac	agcaacaggg	tggtggacct	1020
	catggcctac	atggcctcca	aggagtaaga	aaccctggac	caccaccccc	agcaaggaca	1080
25	ctgagcaaga	gaggccctat	cccaactcgg	cccccaacac	tgagcatctc	cctcacaatt	1140
	tccatcccag	accccataa	taacaggagg	ggcctaggga	gccctcccta	ctctcttgaa	1200
	taccatcaat	aaagttcgct	gcaccac				1228

<210> 4
<211> 333
<212> PRT
<213> Mus musculus

30

<400> 4

35	Met	Val	Lys	Val	Gly	Val	Asn	Gly	Phe	Gly	Arg	Ile	Gly	Arg	Leu	Val
	1				5					10					15	
	Thr	Arg	Ala	Ala	Ile	Cys	Ser	Gly	Lys	Val	Glu	Ile	Val	Ala	Ile	Asn
				20					25					30		
	Asp	Pro	Phe	Ile	Asp	Leu	Asn	Tyr	Met	Val	Tyr	Met	Phe	Gln	Tyr	Asp
40			35				40						45			
	Ser	Thr	His	Gly	Lys	Phe	Asn	Gly	Thr	Val	Lys	Ala	Glu	Asn	Gly	Lys
		50					55					60				
	Leu	Val	Ile	Asn	Gly	Lys	Pro	Ile	Thr	Ile	Phe	Gln	Glu	Arg	Asp	Pro
	65				70					75					80	
45	Thr	Asn	Ile	Lys	Trp	Gly	Glu	Ala	Gly	Ala	Glu	Tyr	Val	Val	Glu	Ser
				85					90						95	
	Thr	Gly	Val	Phe	Thr	Thr	Met	Glu	Lys	Ala	Gly	Ala	His	Leu	Lys	Gly
				100					105					110		
	Gly	Ala	Lys	Arg	Val	Ile	Ile	Ser	Ala	Pro	Ser	Ala	Asp	Ala	Pro	Met
50			115					120					125			
	Phe	Val	Met	Gly	Val	Asn	His	Glu	Lys	Tyr	Asp	Asn	Ser	Leu	Lys	Ile
		130					135					140				
	Val	Ser	Asn	Ala	Ser	Cys	Thr	Thr	Asn	Cys	Leu	Ala	Pro	Leu	Ala	Lys
	145					150					155				160	
55	Val	Ile	His	Asp	Asn	Phe	Gly	Ile	Val	Glu	Gly	Leu	Met	Thr	Thr	Val
				165						170					175	
	His	Ala	Ile	Thr	Ala	Thr	Gln	Lys	Thr	Val	Asp	Gly	Pro	Ser	Gly	Lys
				180					185					190		
	Leu	Trp	Arg	Asp	Gly	Arg	Gly	Ala	Ala	Gln	Asn	Ile	Ile	Pro	Ala	Ser
60			195					200					205			
	Thr	Gly	Ala	Ala	Lys	Ala	Val	Gly	Lys	Val	Ile	Pro	Glu	Leu	Asn	Gly
		210					215					220				

Lys Leu Thr Gly Met Ala Phe Arg Val Pro Thr Pro Asn Val Ser Val
 225 230 235 240
 Val Asp Leu Thr Cys Arg Leu Glu Lys Pro Ala Lys Tyr Asp Asp Ile
 245 250 255
 5 Lys Lys Val Val Lys Gln Ala Ser Glu Gly Pro Leu Lys Gly Ile Leu
 260 265 270
 Gly Tyr Thr Glu Asp Gln Val Val Ser Cys Asp Phe Asn Ser Asn Ser
 275 280 285
 10 His Ser Ser Thr Phe Asp Ala Gly Ala Gly Ile Ala Leu Asn Asp Asn
 290 295 300
 Phe Val Lys Leu Ile Ser Trp Tyr Asp Asn Glu Tyr Gly Tyr Ser Asn
 305 310 315 320
 Arg Val Val Asp Leu Met Ala Tyr Met Ala Ser Lys Glu
 325 330
 15
 <210> 5
 <211> 1705
 <212> DNA
 <213> Mus musculus
 20
 <400> 5
 gccgcgggtga gggaagtgga cgcgatggcc ggggccgcgt ggggtgtccaa ggtctctcgg 60
 ctgctgggtg cattccacaa cacaaaacag gtgacaagag gttttgctgg tgggtgttcag 120
 25 acagtaactt taattccttg agatggaatt ggcccagaaa tttcagcctc agtcatgaag 180
 atttttgatg ctgccaaagc acctattcag tgggaggagc gcaatgtcac agcaattcaa 240
 ggaccaggag gaaagtggat gatccctcca gaagccaagg agtccatgga taagaacaag 300
 atgggcttga aaggcccact aaagacccca atagccgctg gccatccatc tatgaatctg 360
 ttgottogta agacatttga cctttatgcc aatgtccggc catgtgtctc aattgaagggt 420
 tataaaaccc cttacacgga tgtaaatatc gtcaccatcc gagagaacac ggaaggagaa 480
 30 tacagtggaa ttgagcatgt gatcgttgat ggggttgtgc agagcatcaa gctcatcacc 540
 gaagaagcaa gcaagcgcac tgcagagttt gcttcgagta cgctcggaac aaccacogga 600
 accacgtcac ngctgtgcac aaaagctaac atcatgagga tgtcagatgg gctctttctg 660
 caaaaatgca gggaaatttg cggaagaact gtaaagactt aaatttaacg agatgtactt 720
 ggatactgtg gtttaaatat gggtanaaag acccntccaa tttgatgttc ttgtcatgcc 780
 35 aaatttatac ggagacatcc ttagtgatct gtgtgcagga ctgattggag gtcttgggggt 840
 gactccaagt ggcaatattg gagccaacgg tgttgccatc tttgaatcgg ttcattggaac 900
 agccccggac attgcaggca aggacatggc caaccccacg gccctcctgc ttagtgctgt 960
 gatgatgctt cgccacatgg gactttttga ccatgcagca aaaatcgagg ctgcatgttt 1020
 tgctacaatt aaggatggaa agagcttaac aaaagatctg ggaggcaacg cgaagtgtct 1080
 40 tgacttcaca gaagaaatct gtcgtagagt caaagactta gattagcact cctgctgggtg 1140
 gatttgctgc agtcagtcaa tcaactccaaa aggataccct gtaatcctcc ttgagggcgc 1200
 ccaccatttg tttgcttgggt tcttgacaga gtacgttttt tgaatctggc cttttcttaa 1260
 caaaaccctt tgcaatggat gcacatgntg gccccaggcc tttcattcaa aaggtttncc 1320
 ccaagtgtct gtggtattta ttgtcccgtc tgggtaaach ttattttgta aactgtaagt 1380
 45 gaactgtatc atttatcatt gttaacccat tttacacttc aggcaaaatc attttcctca 1440
 actgtaaata ttctgatata gaattaataa gagaagatat ttaacttttt aacaaaagcc 1500
 ctggattttt ggtttatgaa aaacaaactg ggaataaaac agggttttta caatcgaca 1560
 agataacatt attctaatac taatgggtac aaaagaaatt tactgggaaa gttcacagca 1620
 aaaaaatggt atattttctta aaaatatgga aataaagtat ttgtcctata catgaattac 1680
 50 tattaataaaa aatgtaagct ccaag 1705
 <210> 6
 <211> 233
 <212> PRT
 55 <213> Mus musculus
 <400> 6
 Ala Ala Val Arg Glu Val Asp Ala Met Ala Gly Ser Ala Trp Val Ser
 1 5 10 15
 60 Lys Val Ser Arg Leu Leu Gly Ala Phe His Asn Thr Lys Gln Val Thr
 20 25 30
 Arg Gly Phe Ala Gly Gly Val Gln Thr Val Thr Leu Ile Pro Gly Asp

	Gly	Ile	Gly	Pro	Glu	Ile	Ser	Ala	Ser	Val	Met	Lys	Ile	Phe	Asp	Ala	
	50						55					60					
5	Ala	Lys	Ala	Pro	Ile	Gln	Trp	Glu	Glu	Arg	Asn	Val	Thr	Ala	Ile	Gln	
	65				70					75						80	
	Gly	Pro	Gly	Gly	Lys	Trp	Met	Ile	Pro	Pro	Glu	Ala	Lys	Glu	Ser	Met	
					85				90						95		
	Asp	Lys	Asn	Lys	Met	Gly	Leu	Lys	Gly	Pro	Leu	Lys	Thr	Pro	Ile	Ala	
				100					105					110			
10	Ala	Gly	His	Pro	Ser	Met	Asn	Leu	Leu	Leu	Arg	Lys	Thr	Phe	Asp	Leu	
			115					120					125				
	Tyr	Ala	Asn	Val	Arg	Pro	Cys	Val	Ser	Ile	Glu	Gly	Tyr	Lys	Thr	Pro	
	130					135						140					
15	Tyr	Thr	Asp	Val	Asn	Ile	Val	Thr	Ile	Arg	Glu	Asn	Thr	Glu	Gly	Glu	
	145				150						155					160	
	Tyr	Ser	Gly	Ile	Glu	His	Val	Ile	Val	Asp	Gly	Val	Val	Gln	Ser	Ile	
					165					170					175		
	Lys	Leu	Ile	Thr	Glu	Glu	Ala	Ser	Lys	Arg	Ile	Ala	Glu	Phe	Ala	Ser	
				180					185					190			
20	Ser	Thr	Leu	Gly	Thr	Thr	Thr	Gly	Thr	Thr	Ser	Xaa	Leu	Cys	Thr	Lys	
			195					200					205				
	Ala	Asn	Ile	Met	Arg	Met	Ser	Asp	Gly	Leu	Phe	Leu	Gln	Lys	Cys	Arg	
	210						215						220				
25	Glu	Ile	Cys	Gly	Arg	Thr	Val	Lys	Thr								
	225				230												
	<210>	7															
	<211>	853															
	<212>	DNA															
30	<213>	Mus musculus															
	<400>	7															
35	gccatgtttg	gagagagaag	agccaaacag	cctccttcct	gcacagtcc	tcaagctcac											60
	ctcctgcctt	ccgtggacaa	gagggaagcac	aaagaatcat	ccagggtatgg	aagctgaggg											120
	ttccagcgcg	aaggtcacca	ggctaactccg	cctggggagtc	aaggaagact	cggaagaaca											180
	gcatgatgtg	aaagcagagg	ctttcttcca	ggctgggagag	gggagagatg	agcaagggtgc											240
	acaggggccag	cctggagtg	gagcgggtggg	aacagaaggc	gaaggagaag	aattaatgg											300
	aggaaaaggc	cacttttgtc	ctggtgctcc	tggtcctatg	ggtgatgggg	acaaggtag											360
	tggcaccagg	gctggtgggtg	tggagcagga	acaaaatgag	ccagttgctg	agggcactga											420
40	gagccaggag	aatggaaatc	ctgggggtag	gcagatgccc	ctccagggct	ctaggttcgc											480
	ccagcatcga	ctgagggaac	tggagtccat	tttgcagcgc	actaatccct	ttgatgtccc											540
	aaggagggat	cttgateagac	tgateagatgc	ctgtgtgtcc	agagtgcaga	attggtttaa											600
	gatcaggagg	gctgcggcaa	gaagaaccag	gaggagggca	acaccagtcc	ctgaacattt											

Gly Gly Lys Gly His Phe Gly Pro Gly Ala Pro Gly Pro Met Gly Asp
65 70 75 80
Gly Asp Lys Asp Ser Gly Thr Arg Ala Gly Gly Val Glu Gln Glu Gln
85 90 95
5 Asn Glu Pro Val Ala Glu Gly Thr Glu Ser Gln Glu Asn Gly Asn Pro
100 105 110
Gly Gly Arg Gln Met Pro Leu Gln Gly Ser Arg Phe Ala Gln His Arg
115 120 125
10 Leu Arg Glu Leu Glu Ser Ile Leu Gln Arg Thr Asn Ser Phe Asp Val
130 135 140
Pro Arg Glu Asp Leu Asp Arg Leu Met Asp Ala Cys Val Ser Arg Val
145 150 155 160
Gln Asn Trp Phe Lys Ile Arg Arg Ala Ala Ala Arg Arg Asp Arg Arg
165 170 175
15 Arg Ala Thr Pro Val Pro Glu His Phe Arg Gly Thr Phe Glu Cys Pro
180 185 190
Ala Cys Arg Gly Val Arg Trp Gly Glu Arg Cys Pro Phe Ala Thr Pro
195 200 205
20 Arg Phe
210

<210> 9
<211> 882
<212> DNA
25 <213> Mus musculus

<400> 9
ggcacgagct ccggcctagg agaccgcgat ggcgttcctg tgtcagcgcg acagctacgc 60
acgggaggttc accaccaccg tgggtctcctg tagtcccgcc gagctgcaga cggacgcgag 120
30 cggcgggcaag aaagaagtgt tgagcggatt ccatgtggtt ctggaagaca cgctgctttt 180
ccccgagggc gggggacagc ctgatgaccg tggtagcatc aatgacatct ctgtgctgag 240
ggtgacccgt cgtggggccc aggccgatca cttcacggag tcacctctgt cccctgggag 300
tcaagtccag gtccgggtgg actgggagcg gaggtttgac cacatgcagc agcattcagg 360
gcaacatctc atcaccgcgg ttgctgacct tctcttcggg ctgaagacaa cgatcatggga 420
35 gttaggggaga ctccggagtg tgattgagtt ggacagccct tctgtgactg ctgagcaggt 480
ggctgctatc gagcaagagt gtcaatcaga aaatcagaga ccggctgcct ggtgagtgtt 540
cgagagctga nctgnatgac cctgaaggtg gaccaggtga agggggncgg ggntttgccc 600
gatgaatcaa tgctgggccc attcgagntt nttccaatc ggaaggtgtt aaaacnccaa 660
caatgtgcng tngggaacnc acgtnaagcc aaacctcgt gaccttcagg tcnattnaaa 720
40 attctggggg anctggaaaa agggnaaaaa agnacaaaa gccaacctgg atatttcnng 780
gccgggnanc nggttantga antttggatt gggnacnaaa ttcctgggaa gttgaaaaag 840
cctttgnacc ccctaactt naaantntgn gcattnaaan tn 882

<210> 10
45 <211> 171
<212> PRT
<213> Mus musculus

<400> 10
50 Glu Thr Ala Met Ala Phe Leu Cys Gln Arg Asp Ser Tyr Ala Arg Glu
1 5 10 15
Phe Thr Thr Thr Val Val Ser Cys Ser Pro Ala Glu Leu Gln Thr Asp
20 25 30
Ala Ser Gly Gly Lys Lys Glu Val Leu Ser Gly Phe His Val Val Leu
35 40 45
55 Glu Asp Thr Leu Leu Phe Pro Glu Gly Gly Gly Gln Pro Asp Asp Arg
50 55 60
Gly Thr Ile Asn Asp Ile Ser Val Leu Arg Val Thr Arg Arg Gly Ala
65 70 75 80
60 Gln Ala Asp His Phe Thr Glu Ser Pro Leu Ser Pro Gly Ser Gln Val
85 90 95
Gln Val Arg Val Asp Trp Glu Arg Arg Phe Asp His Met Gln Gln His

```

100          105          110
Ser Gly Gln His Leu Ile Thr Ala Val Ala Asp Leu Leu Phe Gly Leu
115          120          125
Lys Thr Thr Ser Trp Glu Leu Gly Arg Leu Arg Ser Val Ile Glu Leu
5      130          135          140
Asp Ser Pro Ser Val Thr Ala Glu Gln Val Ala Ala Ile Glu Gln Glu
145          150          155          160
Cys Gln Ser Glu Asn Gln Arg Pro Ala Ala Trp
165          170

10      <210> 11
      <211> 1464
      <212> DNA
      <213> Mus musculus

15      <400> 11
tcaaggacgc tcaccctggt cttggggcgc caggtttctg ggctgcagcc gcacctggaa      60
aggaggaatc gggagattga tgccagcata aggctggggc tgcaggggaag gcagggttcc      120
atctggtgga ggcagcagag tgaaggcctg cagcagacga gccagcacca caaagagctc      180
cagccgtgcc agaggctctc ccaggcacac gcgtgccccca cagccaaagg atggtgttct      240
gggattcttc ccaggttcca ggaagcgatc tggccagaac ttgctgggca gttcccaaac      300
catctcatcc aggttggcgc cttggatggt ggggatgatg accatatcct tagggatgtc      360
atagccggag atgctgctag ccctagtgtg acgatggggc aaggccaagg gcaccacagg      420
ccgcaaacgc agcacctcgg caatggtggc catgagcaga ggcagctgca ttcggttcct      480
25      gtacaggagc tgggagcctg ggcccagctt gaggtctaac tcttcctgca gtcgcttctg      540
gatctcaggg tggatgaagca ggaaagccac agcccaggag agcgtggtag ccgtgggtctc      600
ggtgccgcgc atgaacaggt ccaccaccga catgtgcacg tgcccctcgt ggagccgctc      660
ttcgtctttg ccatcccttt gcttctccac tccctggagc atgtagtcaa tcatgtcttt      720
ccattggcct gcaaccaggc tgtccttgtg ccgcttcagc tgctgcttga caatatggtc      780
30      ccgactctct tggatctgct tcagcttctg gaggcctggg ttggggagga acctgagaag      840
gggaattatc gtcaagattt ggatggacca gtggttccag gcttgcaaca agtcctggac      900
acagtcgtga agggctctgta ccaacgtgct gtccctgtct ccaaaaagtga ggcaggagat      960
gatactacaa gtgaggaagg agaattcctt atggatggcc acgggggtgc cagcctgggc      1020
tcgcatgcgc tcacagaact cctgggtcag ctgctctatc agaggctcca tggagtctcg      1080
35      catgccagc atcagggctg agcagagag tttcttgtgg gccttccaca tgagagagta      1140
atcccccagg gacagggtcca agtccatctt tccatttagc atatgggggtc ggccagcaaa      1200
gtccacccac ttttggatca aggcctctc aatggttctg ttagaattta gcaccaccac      1260
atcttgcatc cccaagcgga tctgtagat gggcccgagt ttctgagtga ggccaagcag      1320
gtagatggga aggttaggct gtaggaagtg cagaaaaccc ggggccagag gcgggaggtg      1380
40      cagcttccgg agcttccatt ggcccacag ccagcgggtg ccagctagca gcagcaacag      1440
cagcagcagc ccaggtagca gcat      1464

      <210> 12
      <211> 487
      <212> PRT
      <213> Mus musculus

      <400> 12
Met Leu Leu Pro Gly Leu Leu Leu Leu Leu Leu Leu Ala Gly Thr
50      1      5      10      15
Arg Trp Leu Trp Gly Gln Trp Lys Leu Arg Lys Leu His Leu Pro Pro
20      25      30
Leu Ala Pro Gly Phe Leu His Phe Leu Gln Pro Asn Leu Pro Ile Tyr
35      40      45
55      Leu Leu Gly Leu Thr Gln Lys Leu Gly Pro Ile Tyr Arg Ile Arg Leu
50      55      60
Gly Met Gln Asp Val Val Val Leu Asn Ser Asn Arg Thr Ile Glu Glu
65      70      75      80
Ala Leu Ile Gln Lys Trp Val Asp Phe Ala Gly Arg Pro His Met Leu
60      85      90      95
Asn Gly Lys Met Asp Leu Asp Leu Ser Leu Gly Asp Tyr Ser Leu Met
100      105      110

```


Trp Lys Ala His Lys Lys Leu Ser Arg Ser Ala Leu Met Leu Gly Met
115 120 125
Arg Asp Ser Met Glu Pro Leu Ile Glu Gln Leu Thr Gln Glu Phe Cys
130 135 140
5 Glu Arg Met Arg Ala Gln Ala Gly Thr Pro Val Ala Ile His Lys Glu
145 150 155 160
Phe Ser Phe Leu Thr Cys Ser Ile Ile Ser Cys Leu Thr Phe Gly Asp
165 170 175
10 Lys Asp Ser Thr Leu Val Gln Thr Leu His Asp Cys Val Gln Asp Leu
180 185 190
Leu Gln Ala Trp Asn His Trp Ser Ile Gln Ile Leu Thr Ile Ile Pro
195 200 205
Leu Leu Arg Phe Leu Pro Asn Pro Gly Leu Gln Lys Leu Lys Gln Ile
210 215 220
15 Gln Glu Ser Arg Asp His Ile Val Lys Gln Gln Leu Lys Arg His Lys
225 230 235 240
Asp Ser Leu Val Ala Gly Gln Trp Lys Asp Met Ile Asp Tyr Met Leu
245 250 255
20 Gln Gly Val Glu Lys Gln Arg Asp Gly Lys Asp Glu Glu Arg Leu His
260 265 270
Glu Gly His Val His Met Ser Val Val Asp Leu Phe Ile Gly Gly Thr
275 280 285
Glu Thr Thr Ala Thr Thr Leu Ser Trp Ala Val Ala Phe Leu Leu His
290 295 300
25 His Pro Glu Ile Gln Lys Arg Leu Gln Glu Glu Leu Asp Leu Lys Leu
305 310 315 320
Gly Pro Gly Ser Gln Leu Leu Tyr Arg Asn Arg Met Gln Leu Pro Leu
325 330 335
30 Leu Met Ala Thr Ile Ala Glu Val Leu Arg Leu Arg Pro Val Val Pro
340 345 350
Leu Ala Leu Pro His Arg Ala Thr Arg Ala Ser Ser Ile Ser Gly Tyr
355 360 365
Asp Ile Pro Lys Asp Met Val Ile Ile Pro Asn Ile Gln Gly Ala Asn
370 375 380
35 Leu Asp Glu Met Val Trp Glu Leu Pro Ser Lys Phe Trp Pro Asp Arg
385 390 395 400
Phe Leu Glu Pro Gly Lys Asn Pro Arg Thr Pro Ser Phe Gly Cys Gly
405 410 415
40 Ala Arg Val Cys Leu Gly Glu Pro Leu Ala Arg Leu Glu Leu Phe Val
420 425 430
Val Leu Ala Arg Leu Leu Gln Ala Phe Thr Leu Leu Pro Pro Pro Asp
435 440 445
Gly Thr Leu Pro Ser Leu Gln Pro Gln Pro Tyr Ala Gly Ile Asn Leu
450 455 460
45 Pro Ile Pro Pro Phe Gln Val Arg Leu Gln Pro Arg Asn Leu Ala Pro
465 470 475 480
Gln Asp Gln Gly Glu Arg Pro
485

50 <210> 13
<211> 985
<212> DNA
<213> Mus musculus

55 <400> 13
ggcacgagca acagctgaag aggcctccac agaaaccgct gttgaagagt cgggtgaagc 60
tggagaggac ggcgctcctg agggatatggc agagactggg ggaagaggca gcggaagaa 120
ggaggctggg tccagcaaga gcaccgtgga cgggaagctg gtaaaagctt cttttgctcc 180
aataagcttt gccatcaaag ccatgaaaga aggacggtac acggtgctag ctgagagcaa 240
60 gaacgaggag aagaaaaagt ctgggccaac ctctgacaac gaagagggaag atgatgagga 300
agatgggagt tacctgcacc cgtctctctt tgcttccaag aagagcagcc gcctggaaga 360
gctgatgaag cccttgaagg tgggtggacc ggatcacct ctacgagccc tttgtccgga 420

aagcacaaaag ctgacagctc actcccagcc ccaccccaac ttgcaagaat ggggcccgaag 480
 aacgatcaaa gatttgattt ctttcagccc tggcaccag tacaacgcct actacgagtt 540
 ttaaagaagc agttcttcct gcagaaagaa ggaggcggta gcacacaggc agcatcaaca 600
 gctgaagagg ctcccacaga aaccgctgtt gaagagtcgg gaaagcagct cgccatcaac 660
 5 cagacatgtc cgaaggcggg aatccagcca ggagctctcc ttccgcaatg aagacattgg 720
 agtcatggga cctctcagcg gccctgtccg tcctgatcca gcggtaatgg gaacacgcct 780
 ggagctgtcc tgtgattcat tgccccctc cattactgct ctgagcctat tcatgtcagc 840
 accatctgtc taaaatcact acccaggaat gctttttaac gtcgatgatca catctaaatg 900
 aggtcactct gccagcctga ggcttctagg tctgcagagg aaggaagcct tttctctgca 960
 10 gaaaataaaa aggacatgt gcaat 985

<210> 14
 <211> 180
 <212> PRT
 15 <213> Mus musculus

<400> 14
 Ala Arg Ala Thr Ala Glu Glu Ala Pro Thr Glu Thr Ala Val Glu Glu
 1 5 10 15
 20 Ser Gly Glu Ala Gly Glu Asp Gly Ala Pro Glu Gly Met Ala Glu Thr
 20 25 30
 Gly Gly Arg Gly Ser Gly Lys Lys Glu Ala Gly Ser Ser Lys Ser Thr
 35 40 45
 25 Val Asp Gly Lys Leu Val Lys Ala Ser Phe Ala Pro Ile Ser Phe Ala
 50 55 60
 Ile Lys Ala Met Lys Glu Gly Arg Tyr Thr Val Leu Ala Glu Ser Lys
 65 70 75 80
 Asn Glu Glu Lys Lys Lys Ser Gly Pro Thr Ser Asp Asn Glu Glu Glu
 85 90 95
 30 Asp Asp Glu Glu Asp Gly Ser Tyr Leu His Pro Ser Leu Phe Ala Ser
 100 105 110
 Lys Lys Ser Ser Arg Leu Glu Glu Leu Met Lys Pro Leu Lys Val Val
 115 120 125
 35 Asp Pro Asp His Pro Leu Ala Leu Cys Pro Glu Ser Thr Lys Leu
 130 135 140
 Thr Ala His Ser Gln Pro His Pro Asn Leu Gln Glu Trp Gly Pro Arg
 145 150 155 160
 Thr Ile Lys Asp Leu Ile Ser Phe Ser Pro Gly Thr Gln Tyr Asn Ala
 165 170 175
 40 Tyr Tyr Glu Phe
 180

<210> 15
 <211> 612
 45 <212> DNA
 <213> Mus musculus

<400> 15
 cggcacgagc tcgtgtcctc tgcgcctgtg tccgcgcgct gccgcgggcc ttcgcgcgcg 60
 50 tgccccggct tcccacnttg gccttggcgc ggccgctcag caccacccta tgccctgagg 120
 gaatccggag gagaccggg gctctgcagt ccgccttggc gctcgcgcag gtgcctggaa 180
 cagtcacaca tttgtgcgc cagtacagt acgcaccccc actgacgtta gaggaatcaa 240
 ggaccgagtt ctgtatgtct tgaaactcta tgataagatt gatccagaaa agctctccgt 300
 aaattctcat tttatgaagg acctgggctt agacagtttg gaccaagtgg aaattattat 360
 55 ggccatggaa gacgaatttg ggtttgaaat tcctgatata gatgcagaga agttaatgtg 420
 tccacaagaa attgtagatt acattgcaga taagaaggat gtgtatgaat aaagtatcag 480
 agccttcttc ctactgtga ggactccaga ggacacacga tggcatcgtg gccgactgac 540
 agcggctctg ttcaacttgt atttaaattg tctgagtgtt ttacccggtta aaaataaatc 600
 tattacaaaa ct 612

<210> 16
 <211> 86

<212> PRT

<213> Mus musculus

<400> 16

5 Arg Thr Pro Thr Asp Val Arg Gly Ile Lys Asp Arg Val Leu Tyr Val
1 5 10 15
Leu Lys Leu Tyr Asp Lys Ile Asp Pro Glu Lys Leu Ser Val Asn Ser
20 25 30
His Phe Met Lys Asp Leu Gly Leu Asp Ser Leu Asp Gln Val Glu Ile
35 40 45
10 Ile Met Ala Met Glu Asp Glu Phe Gly Phe Glu Ile Pro Asp Ile Asp
50 55 60
Ala Glu Lys Leu Met Cys Pro Gln Glu Ile Val Asp Tyr Ile Ala Asp
65 70 75 80
15 Lys Lys Asp Val Tyr Glu
85

<210> 17

<211> 1036

20 <212> DNA

<213> Mus musculus

<400> 17

25 ggcacgaggg aggagggggc tgggcctgtt gaccgactgc cagtgagagg gaaagctggc 60
aagtttaagg atgatcccga gaagggggca aggtcttccc gctttactag tgtaaacat 120
gatgcgaagg aagagtgtgg caaggtagaa tcaccccctg cagcgagggtg ctcggtcgc 180
agagctgagc tctcgaagca gaatggctcc tcggcctctc agatttcttc tgctgaaggc 240
agggcagctg caaaaggtaa caacagcttg gagagggaga ggcagaattt accaggagcc 300
cttgttctaa acttacaatg aaaccagtca gtcaattaga ctaaagttgt tgattccttg 360
30 tgattatttc catgtgaaaa tggttgtgta caatgacatt taataaaaaat catcctctcg 420
tttagaaggt agaaaggggg gaaaggaaac tttctaaatg ctgcttgaga ttgcagtaag 480
aacatacatt ttctaacctg aaagttgaaa caaatccacac ttgttctgta gactgtgtct 540
ctcttacctg ttgctgtcag gggttaccta tctgctaaac tatgtcggga aagaaaaaat 600
tacttttgtt tgcattgtcat ggggttaatgg tccctgtaat ttggcagtggt gtgtaaaagc 660
35 ttattaaagt tcttcttttg ctttgaccca gaacaatggc atcatttgga tttttgtctg 720
aaatcgtgat accaggtaac tccaaattga tcccttgcac ttgcaacaaa agtattgtgt 780
ttcagtggtt tcacctgtag aaaactagtt ttcactagaa atgctcatca gaacacaaa 840
aaaaaaacca tctttaatag gaataagggt tataattgct tgttgtacag aaatgggtga 900
ctaaagagag agaaacaaag cgtgggaaat ttaaaaaaaa aaccacacaga gaaacaatgg 960
40 taaaaaatga atccaaagag tacgggtgag caagtacaaa tcacctttga gaaaacagaa 1020
actgtcagaa tgggtg 1036

<210> 18

<211> 106

45 <212> PRT

<213> Mus musculus

<400> 18

50 Gly Thr Arg Glu Glu Gly Ala Gly Pro Val Asp Arg Leu Pro Val Arg
1 5 10 15
Gly Lys Ala Gly Lys Phe Lys Asp Asp Pro Glu Lys Gly Ala Arg Ser
20 25 30
Ser Arg Phe Thr Ser Val Asn His Asp Ala Lys Glu Glu Cys Gly Lys
35 40 45
55 Val Glu Ser Pro Pro Ala Ala Arg Cys Ser Ala Arg Arg Ala Glu Leu
50 55 60
Ser Lys Gln Asn Gly Ser Ser Ala Ser Gln Ile Ser Ser Ala Glu Gly
65 70 75 80
Arg Ala Ala Ala Lys Gly Asn Asn Ser Leu Glu Arg Glu Arg Gln Asn
85 90 95
60 Leu Pro Gly Ala Leu Val Leu Asn Leu Gln
100 105

<210> 23

<211> 1895
<212> DNA
<213> Homo sapiens

5 <220>
<221> CDS
<222> (49)...(1677)

<400> 23

10	g t t a g a g g c g g c t t g t g t c c a c g g g a c g c g g g c g g a t o t t c t c c g g c c a t g a g g a a g	57
	Met Arg Lys	
	1	
15	c c a g c c g c t g g c t t c c t t c c c t c a c t c c t g a g g t g c t g c t c c t g c c t	105
	Pro Ala Ala Gly Phe Leu Pro Ser Leu Leu Lys Val Leu Leu Leu Pro	
	5 10 15	
20	c t g g c a c c t g c c g c a g c a g g a t t c g a c t c a g g c c c c a c t c c a g g c	153
	Leu Ala Pro Ala Ala Ala Gln Asp Ser Thr Gln Ala Pro Thr Pro Gly	
	20 25 30 35	
25	a g c c c t c t c t c t c c t a c c g a a t a c g a a c g c t t c t t c g c a c t g c t g a c t	201
	Ser Pro Leu Ser Pro Thr Glu Tyr Glu Arg Phe Phe Ala Leu Leu Thr	
	40 45 50	
30	c c a a c c t g g a a g g c a g a g a c t a c c t g c c g t c t c c g t g c a a c c a c g g c	249
	Pro Thr Trp Lys Ala Glu Thr Thr Cys Arg Leu Arg Ala Thr His Gly	
	55 60 65	
35	t g c c g g a a t c c c a c a c t c g t c a g c t g g a c c a a t a t g a a a a c c a c g g c	297
	Cys Arg Asn Pro Thr Leu Val Gln Leu Asp Gln Tyr Glu Asn His Gly	
	70 75 80	
40	t t a g t g c c c g a t g g t g c t g t c t g c t c c a a c c t c c t a t g c c t c c t g g	345
	Leu Val Pro Asp Gly Ala Val Cys Ser Asn Leu Pro Tyr Ala Ser Trp	
	85 90 95	
45	t t t g a g t c t t t c t g c a g t t c a c t c a c t a c c g t t g c t c c a a c c a c g t c	393
	Phe Glu Ser Phe Cys Gln Phe Thr His Tyr Arg Cys Ser Asn His Val	
	100 105 110 115	
50	t a c t a t g c c a a g a g a g t c c t g t t c c a g c c a g t c t c t a t t c t c a	441
	Tyr Tyr Ala Lys Arg Val Leu Cys Ser Gln Pro Val Ser Ile Leu Ser	
	120 125 130	
55	c c t a a c a c t c t c a a g g a g a t a g a a g c t t c a g c t g a a g t c t c a c c c a c c	489
	Pro Asn Thr Leu Lys Glu Ile Glu Ala Ser Ala Glu Val Ser Pro Thr	
	135 140 145	
60	a c g a t g a c c t c c c a t c t c a c c c a c t t c a c a g t g a c a g a a c c g c a g	537
	Thr Met Thr Ser Pro Ile Ser Pro His Phe Thr Val Thr Glu Arg Gln	
	150 155 160	
65	a c c t t c c a g c c c t g g c c t g a g a g g c t c a g c a a c a a c g t g g a a g a g c t c	585
	Thr Phe Gln Pro Trp Pro Glu Arg Leu Ser Asn Asn Val Glu Glu Leu	
	165 170 175	
70	c t a c a a t c c t c c t t g t c c c t g g g a g g c a g g a g c a a g c g c c a g a g c a c	633
	Leu Gln Ser Ser Leu Ser Leu Gly Gly Gln Glu Gln Ala Pro Glu His	
	180 185 190 195	
75	a a g c a g g a g c a a g g a g t g a g c a c a g g c a g g a g c c g a c a c a a g a a c a c	681

	Lys	Gln	Glu	Gln	Gly	Val	Glu	His	Arg	Gln	Glu	Pro	Thr	Gln	Glu	His	
					200					205					210		
5	aag	cag	gaa	gag	ggg	cag	aaa	cag	gaa	gag	caa	gaa	gag	gaa	cag	gaa	729
	Lys	Gln	Glu	Glu	Gly	Gln	Lys	Gln	Glu	Glu	Gln	Glu	Glu	Glu	Gln	Glu	
				215					220					225			
10	gag	gag	gga	aag	cag	gaa	gaa	gga	cag	ggg	act	aag	gag	gga	cgg	gag	777
	Glu	Glu	Gly	Lys	Gln	Glu	Glu	Gly	Gln	Gly	Thr	Lys	Glu	Gly	Arg	Glu	
			230					235					240				
15	gct	gtg	tct	cag	ctg	cag	aca	gac	tca	gag	ccc	aag	ttt	cac	tct	gaa	825
	Ala	Val	Ser	Gln	Leu	Gln	Thr	Asp	Ser	Glu	Pro	Lys	Phe	His	Ser	Glu	
		245					250					255					
	tct	cta	tct	tct	aac	cct	tcc	tct	ttt	gct	ccc	cgg	gta	cga	gaa	gta	873
	Ser	Leu	Ser	Ser	Asn	Pro	Ser	Ser	Phe	Ala	Pro	Arg	Val	Arg	Glu	Val	
	260					265					270					275	
20	gag	tct	act	cct	atg	ata	atg	gag	aac	atc	cag	gag	ctc	att	cga	tca	921
	Glu	Ser	Thr	Pro	Met	Ile	Met	Glu	Asn	Ile	Gln	Glu	Leu	Ile	Arg	Ser	
					280					285					290		
25	gcc	cag	gaa	ata	gat	gaa	atg	aat	gaa	ata	tat	gat	gag	aac	tcc	tac	969
	Ala	Gln	Glu	Ile	Asp	Glu	Met	Asn	Glu	Ile	Tyr	Asp	Glu	Asn	Ser	Tyr	
				295					300					305			
30	tgg	aga	aac	caa	aac	cct	ggc	agc	ttc	ctg	cag	ctg	ccc	cac	aca	gag	1017
	Trp	Arg	Asn	Gln	Asn	Pro	Gly	Ser	Phe	Leu	Gln	Leu	Pro	His	Thr	Glu	
			310					315					320				
35	gcc	ttg	ctg	gtg	ctg	tgc	tat	tgc	atc	gtg	gag	aat	acc	tgc	atc	ata	1065
	Ala	Leu	Leu	Val	Leu	Cys	Tyr	Ser	Ile	Val	Glu	Asn	Thr	Cys	Ile	Ile	
		325					330					335					
	acc	ccc	aca	gcc	aag	gcc	tgg	aag	tac	atg	gag	gag	gag	atc	ctt	ggc	1113
	Thr	Pro	Thr	Ala	Lys	Ala	Trp	Lys	Tyr	Met	Glu	Glu	Glu	Ile	Leu	Gly	
	340					345					350					355	
40	ttc	ggg	aag	tgc	gtc	tgt	gac	agc	ctt	ggg	cgg	cga	cac	atg	tct	acc	1161
	Phe	Gly	Lys	Ser	Val	Cys	Asp	Ser	Leu	Gly	Arg	Arg	His	Met	Ser	Thr	
					360					365					370		
45	tgt	gcc	ctc	tgt	gac	ttc	tgc	tcc	ttg	aag	ctg	gag	cag	tgc	cac	tca	1209
	Cys	Ala	Leu	Cys	Asp	Phe	Cys	Ser	Leu	Lys	Leu	Glu	Gln	Cys	His	Ser	
				375					380					385			
50	gag	gcc	agc	ctg	cag	cgg	caa	caa	tgc	gac	acc	tcc	cac	aag	act	ccc	1257
	Glu	Ala	Ser	Leu	Gln	Arg	Gln	Gln	Cys	Asp	Thr	Ser	His	Lys	Thr	Pro	
			390					395					400				
55	ttt	gtc	agc	ccc	ttg	ctt	gcc	tcc	cag	agc	ctg	tcc	atc	ggc	aac	cag	1305
	Phe	Val	Ser	Pro	Leu	Leu	Ala	Ser	Gln	Ser	Leu	Ser	Ile	Gly	Asn	Gln	
		405					410					415					
	gta	ggg	tcc	cca	gaa	tca	ggc	cgc	ttt	tac	ggg	ctg	gat	ttg	tac	ggc	1353
	Val	Gly	Ser	Pro	Glu	Ser	Gly	Arg	Phe	Tyr	Gly	Leu	Asp	Leu	Tyr	Gly	
	420					425					430					435	
60	ggg	ctc	cac	atg	gac	ttc	tgg	tgt	gcc	cgg	ctt	gcc	acg	aaa	ggc	tgt	1401
	Gly	Leu	His	Met	Asp	Phe	Trp	Cys	Ala	Arg	Leu	Ala	Thr	Lys	Gly	Cys	
					440					445					450		

GenBank accession number: F01482.1

gaa gat gtc cga gtc tct ggg tgg ctc cag act gag ttc ctt agc ttc 1449
 Glu Asp Val Arg Val Ser Gly Trp Leu Gln Thr Glu Phe Leu Ser Phe
 455 460 465

5 cag gat ggg gat ttc cct acc aag att tgt gac aca gac tat atc cag 1497
 Gln Asp Gly Asp Phe Pro Thr Lys Ile Cys Asp Thr Asp Tyr Ile Gln
 470 475 480

10 tac cca aac tac tgt tcc ttc aaa agc cag cag tgt ctg atg aga aac 1545
 Tyr Pro Asn Tyr Cys Ser Phe Lys Ser Gln Gln Cys Leu Met Arg Asn
 485 490 495

15 cgc aat cgg aag gtg tcc cgc atg aga tgt ctg cag aat gag act tac 1593
 Arg Asn Arg Lys Val Ser Arg Met Arg Cys Leu Gln Asn Glu Thr Tyr
 500 505 510 515

20 agt gcg ctg agc cct ggc aaa agt gag gac gtt gtg ctt cga tgg agc 1641
 Ser Ala Leu Ser Pro Gly Lys Ser Glu Asp Val Val Leu Arg Trp Ser
 520 525 530

cag gag ttc agc acc ttg act cta ggc cag ttc gga tgagctggcg 1687
 Gln Glu Phe Ser Thr Leu Thr Leu Gly Gln Phe Gly
 535 540

25 tctattctgc ccacacccca gcccaacctg cccacgttct ctattgtttt gagaccccat 1747
 tgctttcagg ctgccccttc tgggtctgtt actcggcccc tactcacatt tccttggggtt 1807
 ggagcaacag tccagagag ggccacggtg ggagctgcgc cctccttaaa agatgacttt 1867
 acataaaatg ttgatcttca aaaaaaaaa 1895

30 <210> 24
 <211> 543
 <212> PRT
 <213> Homo sapiens

35 <400> 24
 Met Arg Lys Pro Ala Ala Gly Phe Leu Pro Ser Leu Leu Lys Val Leu
 1 5 10 15
 Leu Leu Pro Leu Ala Pro Ala Ala Ala Gln Asp Ser Thr Gln Ala Pro
 20 25 30
 Thr Pro Gly Ser Pro Leu Ser Pro Thr Glu Tyr Glu Arg Phe Phe Ala
 35 40 45
 Leu Leu Thr Pro Thr Trp Lys Ala Glu Thr Thr Cys Arg Leu Arg Ala
 50 55 60
 Thr His Gly Cys Arg Asn Pro Thr Leu Val Gln Leu Asp Gln Tyr Glu
 65 70 75 80
 Asn His Gly Leu Val Pro Asp Gly Ala Val Cys Ser Asn Leu Pro Tyr
 85 90 95
 Ala Ser Trp Phe Glu Ser Phe Cys Gln Phe Thr His Tyr Arg Cys Ser
 100 105 110
 Asn His Val Tyr Tyr Ala Lys Arg Val Leu Cys Ser Gln Pro Val Ser
 115 120 125
 Ile Leu Ser Pro Asn Thr Leu Lys Glu Ile Glu Ala Ser Ala Glu Val
 130 135 140
 Ser Pro Thr Thr Met Thr Ser Pro Ile Ser Pro His Phe Thr Val Thr
 145 150 155 160
 Glu Arg Gln Thr Phe Gln Pro Trp Pro Glu Arg Leu Ser Asn Asn Val
 165 170 175
 Glu Glu Leu Leu Gln Ser Ser Leu Ser Leu Gly Gly Gln Glu Gln Ala
 180 185 190
 Pro Glu His Lys Gln Glu Gln Gly Val Glu His Arg Gln Glu Pro Thr
 195 200 205

U06003.1: E06003

Gln Glu His Lys Gln Glu Glu Gly Gln Lys Gln Glu Glu Gln Glu Glu
 210 215 220
 Glu Gln Glu Glu Glu Gly Lys Gln Glu Glu Gly Gln Gly Thr Lys Glu
 225 230 235 240
 5 Gly Arg Glu Ala Val Ser Gln Leu Gln Thr Asp Ser Glu Pro Lys Phe
 245 250 255
 His Ser Glu Ser Leu Ser Ser Asn Pro Ser Ser Phe Ala Pro Arg Val
 260 265 270
 10 Arg Glu Val Glu Ser Thr Pro Met Ile Met Glu Asn Ile Gln Glu Leu
 275 280 285
 Ile Arg Ser Ala Gln Glu Ile Asp Glu Met Asn Glu Ile Tyr Asp Glu
 290 295 300
 Asn Ser Tyr Trp Arg Asn Gln Asn Pro Gly Ser Phe Leu Gln Leu Pro
 305 310 315 320
 15 His Thr Glu Ala Leu Val Leu Cys Tyr Ser Ile Val Glu Asn Thr
 325 330 335
 Cys Ile Ile Thr Pro Thr Ala Lys Ala Trp Lys Tyr Met Glu Glu Glu
 340 345 350
 20 Ile Leu Gly Phe Gly Lys Ser Val Cys Asp Ser Leu Gly Arg Arg His
 355 360 365
 Met Ser Thr Cys Ala Leu Cys Asp Phe Cys Ser Leu Lys Leu Glu Gln
 370 375 380
 Cys His Ser Glu Ala Ser Leu Gln Arg Gln Gln Cys Asp Thr Ser His
 385 390 395 400
 25 Lys Thr Pro Phe Val Ser Pro Leu Leu Ala Ser Gln Ser Leu Ser Ile
 405 410 415
 Gly Asn Gln Val Gly Ser Pro Glu Ser Gly Arg Phe Tyr Gly Leu Asp
 420 425 430
 30 Leu Tyr Gly Gly Leu His Met Asp Phe Trp Cys Ala Arg Leu Ala Thr
 435 440 445
 Lys Gly Cys Glu Asp Val Arg Val Ser Gly Trp Leu Gln Thr Glu Phe
 450 455 460
 Leu Ser Phe Gln Asp Gly Asp Phe Pro Thr Lys Ile Cys Asp Thr Asp
 465 470 475 480
 35 Tyr Ile Gln Tyr Pro Asn Tyr Cys Ser Phe Lys Ser Gln Gln Cys Leu
 485 490 495
 Met Arg Asn Arg Asn Arg Lys Val Ser Arg Met Arg Cys Leu Gln Asn
 500 505 510
 40 Glu Thr Tyr Ser Ala Leu Ser Pro Gly Lys Ser Glu Asp Val Val Leu
 515 520 525
 Arg Trp Ser Gln Glu Phe Ser Thr Leu Thr Leu Gly Gln Phe Gly
 530 535 540

45 <210> 25
 <211> 20
 <212> DNA
 <213> Mus musculus

50 <400> 25
 gtggacaaga ggaagcaca 20

55 <210> 26
 <211> 19
 <212> DNA
 <213> Homo sapiens

60 <400> 26
 tctccccatc tcactccac 19

60 <210> 27
 <211> 20
 <212> DNA

	<213> Homo sapiens	
	<400> 27	
5	aaggacaggg gactaaggag	20
	<210> 28	
	<211> 20	
	<212> DNA	
10	<213> Homo sapiens	
	<400> 28	
	ccgtacaaat ccagcccgtg	20
	<210> 29	
15	<211> 20	
	<212> DNA	
	<213> Homo sapiens	
	<400> 29	
20	atgtgagtag gggccgagta	20
	<210> 30	
	<211> 21	
	<212> DNA	
25	<213> Homo sapiens	
	<400> 30	
	ttcctgggct gatcgaatga g	21
30	<210> 31	
	<211> 22	
	<212> DNA	
	<213> Homo sapiens	
35	<400> 31	
	gcaaaagagg aagggttaga ag	22
	<210> 32	
	<211> 20	
40	<212> DNA	
	<213> Homo sapiens	
	<400> 32	
	ccgtggtttt catattggtc	20